

THE GENUS *IRIS* (IRIDACEAE) IN RUSSIA: PHYTOCHEMISTRY, BIOLOGICAL ACTIVITY AND APPLICATION IN TRADITIONAL MEDICINE

© 2023 г. N. B. Alexeeva¹, M. M. Dorofeeva^{2, *}, E. A. Varfolomeeva¹, L. N. Mironova³

¹Komarov Botanical Institute, RAS, Saint-Petersburg, Russia

²Saint Petersburg State Forest Technical University, Saint-Petersburg, Russia

³Botanical Garden-Institute Far Eastern Branch of the Russian Academy of Sciences, Vladivostok, Russia

*e-mail: drofa88@mail.ru

Received December 20, 2022;

Revised February 1, 2023;

Accepted March 14, 2023

The genus *Iris* in Russia is represented by 41 species, of which 4 species are endemic, 11 are included in the Red Book of the Russian Federation and 30 have various regional conservation status. The review provides information on the phytochemical compounds, biological activity and medicinal properties of 16 species growing in Russia, and on their use in traditional medicine of different peoples. The analysis of available data shows that studying the component composition, biological activity levels, and medicinal use of *Iris* L. species requires their correct identification and the knowledge concerning their geographical distribution, ecology, and morphological characteristics. The use of *Iris* species in traditional medicine is described. The extracts and compounds isolated from the most of the studied species exhibit analgesic, antioxidative, antipyretic, anti-inflammatory, antibacterial and antimicrobial activity. Some species also have cytotoxic, antitumor, anti-diabetic, anti-influenza, neuroprotective, antihyperglycemic, antiallergic, antifeedant and other properties. Most of the isolated metabolites were flavonoids, isoflavonoids, anthocyanes, terpenoids, xanthones, quinones, phenolic and fatty acids. In official and traditional medicine in Asia and Europe, the underground and aboveground parts of *Iris aphylla*, *I. lactea*, *I. pseudacorus*, *I. ruthenica*, *I. sanguinea* and other species of the genus are used. The article results from the long-term studies of irises *in situ* and in the collections of two botanical gardens, as well as extensive examination of literature on the component composition and medicinal properties of the studied species.

Keywords: *Iris*, component composition, biological activity, pharmacological properties, traditional medicine

DOI: 10.31857/S003399462301003X, **EDN:** XXHBZX

Depending on the generic and/or specific concepts of different authors, the genus *Iris* L. (Iridaceae) comprises about 260 species [1], considering both bulbiferous and bulbotuberous plants, or about 180 species [2], if only plants with rhizomes and ensiform leaves are considered. Most species of *Iris* s.l. are distributed in Southwest and Central Asia and the Mediterranean region. According to our data [3], there are 41 species of the genus *Iris* in Russia.

Irises are widely used in the traditional medicine of the peoples of Europe and Asia. Many *Iris* species containing such groups as xanthones are a promising source of natural compounds with antiviral activity, and, thus, the development of therapeutic agents based on them is an urgent issue.

Studying the component composition, biologically active substances, and pharmacological properties of *Iris* L. species, the researcher should be familiar with their geographical distribution, ecology, morphological characteristics, and indicate the origin of plant material and who identified it [4–6].

The analysis of 16 species (Table 1), which component composition and medicinal properties were studied, showed that the works of the following authors reflected the taxonomic discrepancy for species from the subgenus *Apogon*: in section *Limniris* – *I. ensata* Thunb., and in section *Haloiris* – *I. biglumis* Vahl., *I. lactea* Pall., *I. oxypetala* Bunge, *I. pallasii* Fisch.

I. ensata Thunb. was described in 1794 by Thunberg [7] from Japan and corresponds to *I. kaempheri* Sieb., and Thunberg's binomial *I. ensata* is a priority and bears no relationship to the continental saline irises [8].

In the Schroeter's summary [9], the description of *I. ensata* (*I. pallasii* Fisch. var. *chinensis* Fisch.) contains information on two different species, namely, *I. ensata* and also *I. oxypetala*, since it is known that the variety *I. pallasii* var. *chinensis* is a synonym of *I. oxypetala* and belongs to the group of halophytic species, as well as *I. lactea*. *I. oxypetala* Bunge was described from the northeast of China [10].

Table 1. The most important species from the genus *Iris* (Russia), their origin, and environmental demands

Section	Species	Origin	Environment
Subgenus <i>Iris</i> (= Pogon)			
<i>Iris</i>	<i>I. aphylla</i>	Russia: south of the European part. Other regions: Central and Eastern Europe	Dryland
	<i>I. pumila</i>	Russia: south of the European part. Other regions: the east of Central and Southern Europe, Moldova, Ukraine, Transcaucasia and Kazakhstan	Dryland
<i>Oncocyclus</i>	<i>I. acutiloba</i>	Russia: Dagestan. Other regions: Azerbaijan, Northern Iran	Dryland
<i>Psammiris</i>	<i>I. humilis</i>	Russia: south of the European part, Eastern and Western Siberia. Other regions: Kazakhstan, Mongolia and China	Dryland
	<i>I. potaninii</i>	Russia: South of Central and Eastern Siberia. Other regions: Mongolia and China	Dryland
Subgenus <i>Apogon</i>			
<i>Haloiris</i>	<i>I. oxypetala</i>	Russia: Khasansky district of the Primorye Territory. Other regions: China, the Korean peninsula and Afghanistan	Dense saline soils
	<i>I. lactea</i>	Russia: Eastern Siberia (Trans-Baikal Territory). Other regions: North-East Kazakhstan, Mongolia and China	Dense saline soils
	<i>I. biglumis</i> *	Russia: Eastern Siberia, west of the Trans-Baikal Territory. Other regions: Northern Mongolia	Dense saline soils
	<i>I. pallasii</i> *	Russia: Western Siberia. Other regions: North-East Kazakhstan and Mongolia	Dense saline soils
<i>Tenuifolia</i>	<i>I. tenuifolia</i>	Russia: south-east of the European part of Russia, south-western Siberia, the Trans-Baikal Territory. Other regions: Kazakhstan, north-western China and northern and north-eastern Mongolia	Very dry, neutral
<i>Xyridion</i>	<i>I. halophila</i>	Russia: South of the European part of Russia, Krasnodar and Stavropol Territories, Chechnya, Ingushetia, Dagestan and south of Western Siberia. Other regions: Central Europe, Kazakhstan, Mongolia and Afghanistan	Dryland
<i>Limniris</i>	<i>I. ensata</i>	Russia: Yakutia, the Amur Region, South-West of the Khabarovsk Territory, Primorye Territory and the Kuril Islands. Other regions: north-eastern China, the Korean peninsula and Japan	Wetland
	<i>I. laevigata</i>	Russia: Eastern Siberia, Primorye Territory and the Kuril Islands. Other regions: Eastern Mongolia, Northeast China and Japan	Wetland
	<i>I. pseudacorus</i>	Russia: The European part of Russia, Caucasus, Western Siberia. Other regions: Europe, Northern Africa and Asia Minor	Wetland
	<i>I. sanguinea</i>	Russia: Eastern Siberia and the Primorye Territory. Other regions: Mongolia, north-eastern China, the Korean Peninsula and Japan	Wet, moist, or normal
	<i>I. sibirica</i>	Russia: Eastern Europe, the Caucasus, Western and Central Siberia. Other regions: Armenia, Kazakhstan	Wet, moist, or normal
<i>Tripetalae</i>	<i>I. setosa</i>	Russia: Central and Eastern Siberia, the Far East. Other regions: Japan, north-eastern China, the Korean Peninsula, the Pacific coast of North America	From moist to wet
<i>Ioniris</i>	<i>I. ruthenica</i>	Russia: south of Siberia. Other regions: in Asia — Kazakhstan, north-western Mongolia and China; in Europe — the Romanian Carpathians	Slightly dry

* No data on the component composition and biological activity.

Phytochemical studies of the aboveground part of *I. ensata* auc. non. Thunb. [11–16] at Leningrad Chemical and Pharmaceutical Institute (now St. Petersburg State University of Chemical Pharmacy) should be attributed to *I. lactea*. Pall. [17].

In “Plant Resources...” reference book [18, 19], description of *I. ensata* provides information on two different species: *I. ensata* and *I. lactea*, as the distribution of *I. ensata* is indicated in the meadows of the Kuril Islands (Kunashir Island) on saline soils, and in steppes, though there are no steppes on Kunashir Island. Other species from the group of halophytic irises are listed under *I. biglumis* s.l. The taxonomic divergence is also found in the works of modern authors [20].

To avoid errors, in our review 4 saline species of the sect. *Haloiris*, are jointly referred to as *I. lactea* s.l. All works referring to the sect. *Haloiris* are grouped as follows:

I. oxypetala Bunge is mentioned in publications under the following names: *I. lactea* var. *chinensis* (Fisch.) Koidz., *I. pallasii* Fisch. var. *chinensis* Fisch., and *I. lactea pallasii* Fischer var. *chinensis* [21];

I. lactea Pall. is mentioned in publications as *I. lactea* Pall. var. *lactea*, *I. oxypetala* C.A. Mey, and *I. ensata* Thunb.) [22].

Representatives of the sect. *Haloiris* are widely used in traditional medicine in Russia, China, Mongolia, India and in Tibetan medicine. In traditional medical practice, all parts of *I. lactea* s.l. are utilized. Rhizomes and seeds are used for the treatment of gastroenteritis, hepatitis, fever, kidney disease, and as anti-inflammatory and antixudative agents. Flowers are used for the treatment of gastrointestinal and gallstone diseases, fruits – for wounds, ulcers and burns. All parts of plants are anthelmintic, and produce immunostimulant, anti-inflammatory, cardiogenic and nephroprotective effects [23–27]. Seeds contain compounds that, among others, exhibit anti-inflammatory, antioxidant, antitumor, and antiradiation effects and are used in the treatment of lung cancer, esophageal cancer, head and neck cancer as an antineoplastic agent and radiosensitizer [28–32].

Also, all parts of *I. lactea* s.l. are used in traditional medicine for pneumonia, bronchitis, chronic gastritis and jaundice [24]. Pills or powder, or 3–9 g of decoction of *I. pallasii* var. *chinensis* are used for jaundice, nosebleeds, hematemesis, ulcers and boils, diarrhea, leucorrhoea, pharyngitis, metrorrhagia [33].

Resulting from extensive pharmacological studies carried out at St. Petersburg Chemical and Pharmaceutical Academy, Laktir, a drug in the form of tablets was developed. It has a pronounced anti-inflammatory effect at different stages of inflammation [34].

All species in the sect. *Limniris* have pronounced pharmacological properties. *I. pseudacorus* L. is commonly used in traditional medicine of Europe and Asia, due to its wide distribution in Russia and other countries. Its distribution range covers vast areas from

the middle of taiga to the subtropical zone in Europe, the Caucasus, Western Siberia, North Africa and Asia Minor, as mentioned in numerous publications [5, 9, 18–20, 27, 34, 35].

Rhizomes of *I. pseudacorus* are used in the Tibetan and Chinese medicine as an antipyretic agent. Rhizomes, seeds and flowers are used in the treatment of pneumonia, bronchitis, chronic gastritis, jaundice and other liver diseases. All plant parts, especially the fruit, are considered laxative, anthelmintic, hypnotic, and astringent. Seeds are used as hemostatic agent in gynaecology, for the treatment of wounds, injuries, burns, malignant tumors, carbuncles. In traditional medicine a decoction of a fresh rhizome of *I. pseudacorus* is used as an astringent, tonic, laxative, anti-pneumonia, and anti-stomach ulcer agents, a treatment for urinary diseases, ascites, metrorrhagia, headache and toothache, and as hair growth stimulant. Furthermore, the essence is used for epilepsy, and as topical medication for diathesis, stomatitis, and toothache. It is one of the ingredients of M.N. Zdenko polyherbal tea, used in anticancer therapy [9, 18, 19, 36–39].

All parts of *I. sanguinea* Donn. and *I. sibirica* L. are used in traditional medicine of Russia and China. A decoction of *I. sanguinea* rhizomes is used for gynecological diseases, has an emetic, laxative, anthelmintic, and antiedemic action, and as topical medication it exhibits hemostatic effect, and is used in wounds, burns and snake bites healing [24, 40–43]. *I. sibirica* is used as an emetic, laxative, pain reliever, and in the treatment of syphilis, gonorrhea and gynecological diseases [24, 40, 41, 44].

In different countries *I. setosa* Pall. is a popular medicinal plant. In Japanese medicine rhizomes of *I. setosa* are used for scabies, a decoction is used for pneumonia, ascites, tonsillitis; and externally – for wounds, ulcers, fistulas, and as antifreckle treatment; in Yakutia, rhizomes are used as an analgesic (for toothache) and emetic, and flowers – to treat skin abscesses [45, 46]. Rhizomes, flowers and seeds of *I. ruthenica* Ker-Gawl. are used as anthelmintic in Tibetan medicine [47].

In the literature the use of the species in the subgenera *Iris* (= *Pogoniris*) sect. *Iris* and *Psammiris* in traditional medicine is documented. An extract of *I. aphylla* L. exhibit antibacterial, antioxidant, anti-inflammatory and antiallergic activity [48–50]. Rhizomes and leaves of *I. pumila* L. exert antifungal and detoxifying activity [51, 52]. Rhizomes, seeds and flowers of *I. humilis* Georgii are used in Tibetan medicine: the rhizomes reduce intoxication, sepsis, various infectious diseases, and are used as an external hemostatic agent [53, 54]. In Mongolian traditional medicine *I. potaninii* Maxim. is used for the treatment of various diseases, such as bacterial infections, cancer, and inflammation [4].

Table 2. Component composition and biological activity of species *Iris* L.

Species	Plant part	Compounds	Biological activity and use	Ref.
<i>Habiris</i> Doronkin				
<i>I. lactea</i> Pall. (= <i>I. lactea</i> Pull. var. <i>lactea</i> , <i>I. oxypetala</i> C.A. Mey, <i>I. ensata</i> auc. non. Thunb.)	Rhizomes Roots	Xanthones: iriflophenon, iriflophenone 2-<i>O</i>-hexoside, iriflophenone 4-<i>O</i>-hexoside, isomangiferin, mangiferin, nigransiden, 4-<i>O</i>-methyliriflophenon, 7-<i>O</i>-methylisomangiferin, 7-<i>O</i>-methylmangiferin, bellidifolin [73]. Phenolic acids: vanillic, <i>trans</i>-cinnamic, salicylic, <i>p</i>-hydroxybenzoic, protocatechuic, ferulic, caffeic [73]. Phenols: 3-hydroxy-5-methoxyacetophenone, apocynin [73]. Flavonoids and isoflavonoids: apigenin 8-<i>C</i>-(2''-pentosyl)-hexoside, apigenin-6,8-di-<i>C</i>-arabinoxide, luteolin 7-<i>O</i>-(2''-<i>p</i>-coumaroyl)-rhamnoside, iriflophenone 4-<i>O</i>-(6''-acetyl)-hexoside, 4-methyltectorigenin-7-glucoside, 6,4'-dimethoxy-5-hydroxyflavone-7-glucoside, iridin, irilin A, irilin B, irilin D, iristectorigenin A, iristectorigenin A 7-<i>O</i>-hexuronide, iristectorigenin B, irigenin, irigenin S, iristectorin A, iristectorin B, iristectoridin B, irisdichotin A, irisdichotin C, irilone A, irilone B, irilone 4'-<i>O</i>-hexoside, irilone 4'-<i>O</i>-β-d-glucopyranoside, irilone 4'-<i>O</i>-(6''-hexosyl)-hexoside, irilone 4'-<i>O</i>-β-d-glucopyranoside-(2→1)-l-rhamnoside, irilone 4'-<i>O</i>-(6''-(3-hydroxy-3-methylglutaryl))-hexoside, irisolidone, irisolidone 7-<i>O</i>-hexoside, irisolidone-7-<i>O</i>-α-<i>D</i>-glucoside, irisflorentin a, irisflorentin b, irifloside, irisflogenin-4'-<i>O</i>-(β-<i>D</i>-glucopyranosyl)-(1→6)β-<i>D</i>-glucopyranoside], iriskumaonin methyl ether, 6,3',4'-trimethoxy-7,8,5'-trihydroxyisoflavone, 5-methoxy-4'-hydroxy-6,7-methylenedioxyisoflavone, 4',7-di-<i>O</i>-methylidihydroquercetin-5,3',3'-trihydroxy-7,4'-dimethoxyflavanone, 7,4'-dimethoxy-8,3',5'-trihydroxy-6-<i>O</i>-β-<i>D</i>-glucopyranosylisoflavon, dichotomin, genistein, germanism B, isoschaftoside, nigricin, schaftoside, swertiajaponin, tectoridin, tectorigenin [73]. Terpenoids: α-dehydroirigermanal, 22,23-epoxy-10-deoxy-21-hydroxyiridal, 22,23-epoxy-iridal, isoiridogermanal, belamcandal, iridotectoral A, iridotectoral B, iridal, α-irigermanal, amorphin, α-murolene, β-gujunene, γ-elemene, iriversical [73]. Fatty acids: myristic, lauric, palmitic, stearic [73]. Steroids: stigmasterol, stigmasterol-3-<i>O</i>-b-<i>D</i>-glucopyranoside, 7-β-hydroxystigmast-4-en-3-one [73]. Quinones: irisoquin B [73].	<i>In Mongolian medicine</i>, the plant was used as an anthelmintic agent; in <i>Transbaikalia</i> – for infectious and viral diseases [25]. <i>In Tibetan medicine</i> – for kidney diseases (rhizomes), gastrointestinal diseases (flowers), and wounds (topical ointment from fruits) [27]. <i>In China</i> – for cough and cold, liver disorders, seeds and flowers – as antipyretic and detoxifier [77], as a hemostatic and anticolic agent [78]. <i>In India</i> <i>I. lactea</i> is known as “Teshmentok” and whole plant is used as green fodder supplement increasing milk production in cattle [79]. <i>In Trans-Himalayan Region</i> <i>I. lactea</i> is known as “Dres-ma”, whole plant is dried and powdered, <i>decoction</i> is taken orally to improve appetite, as stomach and small and large intestinal cramps reliever, and as food-poisoning remedy [80]. The <i>dry extract</i> of <i>I. lactea herb</i> is nontoxic and produces immunostimulating, anti-inflammatory, cardiotoxic and nephroprotective effects. Extract has an antioxidant effect in toxic hepatitis [26, 34, 81–87]. <i>Dry extract of the rhizomes and roots</i> has anti-inflammatory and antitoxidative properties in pulmonary edema [88]. <i>Methanol extracts from leaves and rhizomes</i> inhibits the adhesion of gram-positive (<i>Staphylococcus aureus</i>) and gram-negative (<i>Pseudomonas aeruginosa</i>) bacteria as well as the dental plaque multispecies biofilm [73]. <i>Embinin</i> from the <i>aqueous extract</i> of <i>I. lactea</i> aerial parts has an inhibitory effect on the proliferative activity of lymphocytes [89]. <i>Embinin</i> exhibits high anti-influenza activity [25]. A technology for obtaining tincture and syrup with <i>I. lactea</i> extract was developed [90, 91].	11, 12, 13, 14, 15, 16, 17, 25, 26, 27, 34, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91
	Aerial parts	Xanthones: mangiferin, isomangiferin [13, 14, 25, 26]. Phenolic acids: ferulic, <i>n</i>-coumaric, vanillic, <i>n</i>-hydroxybenzoic [15], caffeic, chlorogenic, neochlorogenic [25], protocatechuic, syringic, gallic, <i>trans</i>-cinnamic [74]. Flavonoids: 6-<i>C</i>-β-<i>D</i>-glucopyranosyl-<i>O</i>-<i>L</i>-rhamnoside, apigenin 7,4'-dimethyl ether [14], orientin, homoorientin (luteolin-6-<i>C</i>-glucoside, isoorientin) [16], luteolin, apigenin, embinin, 4''-acetylbembinin, 2'',4''-diacetylbembinin [25], swertiajaponin [26, 75]. C-glycosylflavones: diacetylbembinin, acetylbembinin, embigenin [17], 4''-<i>O</i>-acetyl-embinin, 2'',4''-<i>O</i>-diacetyl-embinin, 6'',4''-<i>O</i>-diacetyl-embinin [76].		

Table 2. (Contd.)

Species	Plant part	Compounds	Biological activity and use	Ref.
<i>I. oxypetal</i> Bunge (= <i>I. lactea</i> var. <i>chinesis</i> (Fisch.) Koidz., <i>I. lactea pallasi</i> Fischer var. <i>chinesis</i> , <i>I. pallasi</i> Fisch. var. <i>chinesis</i> Fisch.)	Leaves	Xanthones: iriflophenon, iriflophenone 2- <i>O</i> -hexoside, iriflophenone 4- <i>O</i> -hexoside, isomangiferin, mangiferin, nigricansiden, bellidifolin, 4- <i>O</i> -methyliriflophenon [73]. Phenolic acids: vanillic, <i>trans</i> -cinnamic, salicylic, <i>p</i> -hydroxybenzoic, protocatechuic, ferulic, caffeic [73]. Flavonoids and isoflavonoids: luteolin 7- <i>O</i> -(2''- <i>p</i> -coumaroyl)-rhamnoside, luteolin 6- <i>C</i> -glucoside, luteolin 8- <i>C</i> -hexoside, apigenin 8- <i>C</i> -(2''-hexosyl)-hexoside, apigenin 6- <i>C</i> -hexoside, apigenin 8- <i>C</i> -glucoside, kaempferol 7- <i>O</i> -(6''-rhamnosyl)-hexoside, kaempferol 3- <i>O</i> -galactoside, kaempferol 3- <i>O</i> -glucoside, isoorientin, irilone A, irilin A, irilin B, irilin D, irisolidone, iristectorigenin A, iristectorigenin B, isovitexin, tectorigenin [73]. Terpenoids: α -dehydroirigermanol, 22,23-epoxy-10-deoxy-21-hydroxyiridal, 22,23-epoxyiridal, isoiridogermanol, belamcandal, iridotectoral A, iridotectoral B, spirobicyclic triterpenoid, amorphin, α -muurolene, β -gurjulenene, γ -elemene, iridal, α -irigermanol, irigermanic C [73]. Fatty acids: myristic, lauric, stearic, palmitic [73]. Steroids: stigmastrol-3- <i>O</i> - β - <i>D</i> -glucopyranoside, 7- β -hydroxystigmast-4-en-3-one [73]. Quinones: irisquinone A, irisquinone B, pallasone B, dihydroirisquinone [73]. Contain some traces of alkaloids, show unclear result for saponins and coumarins [11, 12].		
	Seeds	Phenol derivative: 3-[(<i>Z</i>)-10-heptadecenyl]-4,5-dimethoxyphenol (irisphenol), 3-[(<i>Z</i>)-10-heptadecenyl]-5-methoxyphenol [92]. Benzene derivatives: 3,5-dimethoxy-[(<i>Z</i>)-10-heptadecenyl]benzene, 2-[(<i>Z</i>)-10-heptadecenyl]-4,6-dimethoxyphenol (belamcandaphenol P), 1,2,3,4-tetramethylbenzene, pentamethylbenzene, 1-ethyl-2,4,5-trimethylbenzene [92]. Benzofuran derivatives: 9,9b-di[(<i>Z</i>)-10-heptadecenyl]-4a,8-dihydroxy-2,7-dimethoxy-1,4-dioxo-1,4,4a,9b-tetrahydrodibenzofuran (belamcandone P) [93]. Quinones: pallasone B (dihydroirisquinone), pallasone C [94, 95], irisquinone [29, 31, 94, 96], 3-hydroxyirisquinone [92]. Fatty oils content 12%. Higher fatty acids (%) : decanoic (0.02–0.69), lauric (0.02–0.37), tetradecanoic (myristic) (0.08–1.12), stearic (0.89–36), palmitic (5–8.56), oleic (28.51–37.53), linoleic (41.31–65.35) [33], palmitoleic (0.13), arachidonic (1.72), linolenic (0.26), 11-eicosenoic (0.72), docosanoic (0.40), tricosanoic (0.11), lignoceric (0.24), docosahexaenoic (3.18) [92]. Flavan and proanthocyanidin: epicatechin-(4 α →8)-catechin (procyanidin B1), catechin-(4 α →8)-catechin (procyanidin B3), galloocatechin-(4 α →8)-catechin (prodelphinidin B3) [97], catechin, epicatechin-(4 β →6)-catechin (procyanidin B7) [98, 99], 3,4-dihydroxybenzoic acid, 3'- <i>O</i> -(1-hydroxy-6-oxo-2-cyclohexene-1-carboxylic acid ester) of procyanidin B1, 6-[(1S)-3-methoxy-3-oxo-1-(2,4,5-trihydroxyphenyl)propyl]catechin, fisetinidol (4 α ,8)-catechin, epiafzelechin-(4 β →8)-epicatechin, epicatechin, procyanidin A1, nora-thyriol, proanthocyanidin A1, procyanidin B6, (2'S,3'R)-9-(5',6'-dihydroxy-2'-hydroxymethyl)-2',3'-dihydrobenzo[b]furan-3-yloxy)-6H-dibenzo[b,d]pyran-6-oe, hopeaphenol, isohopeaphenol, n-butyl pro-lithospermate [100]. Oligostilbenes: vitisin A, viniferin, <i>e</i> -viniferin, vitisin B, vitisin C [100–104], vitisin D, ampelopsin B, <i>cis</i> -vitisin A [105], <i>trans</i> - <i>e</i> -viniferin, vatalbinoside C, <i>cis</i> - <i>e</i> -viniferin-11a,13b- <i>O</i> - β -d-digluco- <i>trans</i> - <i>e</i> -viniferin-13b- <i>O</i> - β -d-glucopyranose, <i>cis</i> -vitisin B-13b- <i>O</i> - β -d-glucopyranoside, <i>cis</i> -vitisin B, <i>cis</i> -vitisin C [99], vitisin A-13- <i>O</i> - β -d-glucoside, hopeaphenol [106].	In Chinese medicine the seeds are known as Ma Lin Zi and used for cold, phlegm, intestinal wind [92], jaundice, strangury, urine cloudiness, inhibited urination (dysuria), intestinal abscess, intestinal worms accumulation, malaria, pharyngitis, toothache, bloody spotting, spontaneous external bleeding, hematochezia, metrorrhagia and metrorrhagia, swelling of sores, scrofula, hernia, hemorrhoids, scalds, snake bites [95]. Also, seeds are used as treatment for malignancies, exhibit blood activating and detoxifying effect, and can be used for the treatment of metrorrhagia and vaginal discharge [28–30]. In Chinese traditional medicine, the dried seeds of <i>I. lactea</i> var. <i>chinesis</i> are used as treatment for jaundice, diarrhoea, leucorrhoea, pharyngitis, inflammation and carbuncles [108]. In traditional medicine seeds are used as a laxative [24, 44], anthelmintic [24, 47], hypnotic [111], astringent [45], tonic and blood purifier agents, and in therapy of syphilis [44]. Also, seeds are used for gynecological diseases [24], as a hemostatic agent [44, 45, 112], for the treatment of wounds [113], injuries [112], burns [23], malignant tumors [113], carbuncles, as a diuretic [44, 113], for dropsy [44, 114], stranguria, jaundice, and as an antidote to poisonous insect and snake bites [112]. Seeds contain compounds, which exhibit various biological effects, including anti-inflammatory, antioxidant, antitumor, and antiradiation ones [32]. Irisquinone capsules extracted from the seed coat have been registered in the Chinese New Drug Conversion Standard and are used as radiosensitizer. Irisquinone isolated from seeds , has been successfully used in lung, esophageal, head and neck cancer therapy as an antineoplastic agent and radiosensitizer [31]. It is effective against transplantable tumors, such as	11, 12, 23, 24, 28, 29, 30, 31, 32, 33, 42, 44, 45, 47, 62, 71, 92, 93, 94, 95, 96, 97, 98, 99, 100, 101, 102, 103, 104, 105, 106, 107, 108, 109, 110, 111, 112, 113, 114, 115,

Table 2. (Contd.)

Species	Plant part	Compounds	Biological activity and use	Ref.
		Carboxylic acids: acetic acid, hexanoic acid, nonanoic acid. Aldehydes: hexanal, furfural, 2-heptenal, octanal, 5-methylfurfural, 2-octenal, 4-decenal, 1-nonanal, trans-2-nonenal, 5-hydroxymethylfurfural, vanillin. Aromatic alcohol: benzyl alcohol. Fatty alcohols: 1-hexanol, 1-octen-3-ol, 1-octanol, 1-nonen-4-ol, 1-nonanol, 4-phenyl-2-butanol. Heterocyclic compounds: 3-ethyl-2,5-dimethylpyrazine, 4-methylpyridine, 2-furanmethanol, 2-pentylfuran. Terpenes: limonene, <i>trans</i>-geranylgeraniol, squalene, <i>trans</i>-squalene, cycloartan-24-en-1α,2α,3β-triol, lupeol, 2,6-dimethylocta-2,6-diene, (3β,24E)-3-hydroxy-9,19-cyclolanost-24-methylene. Cyclic ether: 2,2-dimethyl-3-(3,7,12,16,20-pentamethyl-3,7,11,15,19-twenty-monopentenyl)-ethylene oxide. Alkanes: dodecane, pentadecane, tridecane, tetradecane, hexadecane, 2,6,10,14-tetramethylpentadecane, <i>n</i>-eicosane, 3-methyltricosane, <i>n</i>-tetracosane, <i>n</i>-pentacosane, 3-methylpentacosane, <i>n</i>-hexacosane, <i>n</i>-heptacosane, <i>n</i>-octacosane, <i>n</i>-nonacosane. Anhydrides: 3-dodecyl-2,5-furandione. Phosphoric esters: tri-<i>n</i>-butyl phosphate. Carboxylic ester: octyl octanoate. Fatty acid derivative: erucamide. Alkenes: 3-heptylacrolein. Aromatic hydrocarbons: naphthalene, benzothiazole, 2-methylnaphthalene, 1-methylnaphthalene, 1,7-dimethylnaphthalene, 1,5-dimethylnaphthalene, 1,2-dimethylnaphthalene, 1-ethylnaphthalene, γ-nonalactone, 1,2-dimethylindan. Aliphatic hydrocarbons: 1-hexacosene. Kenones: 4,7-dimethoxy-1-indanone, benzylacetone. Sterols: campesterol, stigmasterol, β-sitosterol, delta5-avenasterol, betulinicaldehyde, γ-sitosterone [92].	cervical carcinoma U₁₄, lymphosarcoma, hepatoma and Ehrlich ascites carcinoma in mice by peritoneal injection and oral administration. It is also a sensitizer in radiation therapy of cancer [28–30]. The radiosensitizing effect of <i>irisquinone A</i> was detected <i>in vitro</i> against the tumor cell lines U₁₄, S-180V, HeLa, on the mice breast cancer cell lines Ma 7373, and against human intestinal mucadenocarcinoma in hairless mice [28, 115, 116]. The mechanism of this effect was determined to an inhibition of oxygen consumption and depletion of glutathione in tumor cells [116]. In a clinical trial of 558 patients with lung cancer and nasopharyngeal cancer or superficial metastatic cancer during radiotherapy, oral <i>irisquinone A</i> significantly reduced tumor size and prolonged patients' survival [28, 95, 117]. Flavan-3-ol compounds from the <i>seed coat</i> have potential therapeutic benefits in the treatment of human osteoarthritis and for enhancing prostate cancer cell antagonists [101, 118]. Oligostilbenes extracts from the <i>seeds</i> showed potent anti-inflammation activities, exhibited antidiabetic and antilipogenic effects, attenuated hyperglycemia, hyperlipidemia, ameliorated lipid metabolism and attenuated hepatic steatosis in mice [100, 104, 106]. In addition, <i>oligostilbenes</i> isolated from <i>seeds</i> exhibited different bioactivities on adipocytes differentiation and can be useful for the treatment of obesity and obesity-related metabolic diseases. [103]. Vitisin B was a fatty acid synthase inhibitor and exhibited apoptotic effect on human breast cancer cells [102]. In traditional medicine rhizomes, seeds and flowers are used for pneumonia, bronchitis, chronic gastritis [24] and liver diseases [44, 114]. In Tibetan medicine, rhizomes are used as an antipyretic remedy [42, 113]. In Chinese traditional medicine, flowers and roots are used for the treatment of hemorrhoids, ulcer, blood vomiting and stranguria with cloudy discharge, and as an antipyretic agent, the leaves are used as a treatment for pharyngitis and pain in joints of the lower back and legs [119]. Embinin A and irislactin C isolated from the <i>leaves</i> showed weak cytotoxicity against A549 (human lung cancer) cells [109].	116, 117, 118, 119
	Rhizomes Roots	Flavonoids: 5-hydroxy-7-methoxyflavone (tectochrysin), 5,2'-dihydroxy-6,7-methylenedioxyflavanone, irisoid A, irisoid D [62, 107]. Isoflavones: 5,7-dihydroxy-6,2'-dimethoxyisoflavone, 4',5-dihydroxy-6,7-dimethoxyisoflavone, 4',5,7-trihydroxy-6-methoxyisoflavone (tectorigenin), 5,3'-dihydroxy-2'-methoxy-6,7-methylenedioxyisoflavone, 5,2'-dihydroxy-6,7-methylenedioxyisoflavone (irisone B) [62, 107]. Fatty acids: heptadecanoic acid [107]. Contain any traces of alkaloids, shown positive result for flavonoids and coumarins [11, 12].		

Table 2. (Contd.)

Species	Plant part	Compounds	Biological activity and use	Ref.
<i>I. tenuifolia</i> Pall.	Leaves	Flavonoids and isoflavonoid: apigenin 7- <i>O</i> -glucoside-6- <i>C</i> -glucoside, luteolin 6- <i>C</i> - β - <i>D</i> -glucoside, saponaretin, scoparin [32]. C-glycosylflavones: 7- <i>O</i> -(2,4-di- <i>O</i> -acetyl- β - <i>D</i> -Rhap)-6- <i>C</i> -(6- <i>O</i> -acetyl- α -L-Glcp)-(1 $\rightarrow$ 2)- β -D-Glep (irisilactin A), 6- <i>C</i> -(2,3-di- <i>O</i> -acetyl- α -L-Rhap)-(1 $\rightarrow$ 2)- β - <i>D</i> -Glep (irisilactin B) [108], 5-hydroxy-7,4'-dimethoxyflavone-6- <i>C</i> -[<i>O</i> -(α -L-4'''-acetylthamnopyranosyl)-(1 $\rightarrow$ 2)- β - <i>D</i> -glucopyranoside] (embinin A), 5-hydroxy-7,4'-dimethoxyflavone-6- <i>C</i> -[<i>O</i> -(α -L-2'',4'''-diacetylthamnopyranosyl)-(1 $\rightarrow$ 2)- β - <i>D</i> -glucopyranoside] (embinin B), 5-hydroxy-7,4'-dimethoxyflavone-6- <i>C</i> -[<i>O</i> -(α -L-3'',4'''-diacetylthamnopyranosyl)-(1 $\rightarrow$ 2)- β - <i>D</i> -glucopyranoside] (embinin C), 5-hydroxy-4'-methoxyflavone-7- <i>O</i> -(β - <i>D</i> -4'''-acetylthamnopyranosyl)-6- <i>C</i> -[<i>O</i> -(α -L-6'''-acetylglucopyranosyl)-(1 $\rightarrow$ 2)- β - <i>D</i> -glucopyranoside] (irisilactin C) [109], swertiajaponin, swertisin 2''- <i>O</i> -rhamnoside-4'- <i>O</i> -glucoside, swertisin 2''- <i>O</i> -(4'''-acetylthamnosside)-4'- <i>O</i> -glucoside, embinin, 2'''-acetyl-embinin, 3'''-acetyl-embinin, the isomer of irisilactin A, the isomer of irisilactin C [32]. Xanthones: neomangiferin, mangiferin, isomangiferin [32, 71].		
	Flowers	Flavonoids: quercetin, kaempferol, isorhamntin [110].		
		<i>Tenuifolia</i> (Diels) Alexeeva		
Whole plant	Flavonoids: tenuifone, izalpinin, alpinone, irilin B, irisone A, irisone B, betavulgarin, 5,7-dihydroxy-2',6'-dimethoxyisoflavone, 2',5'-dihydroxy-6,7'-methylenedioxy flavanone. Benzene derivatives: tenuifodione. Polynuclear aromatic compounds: arborinone. Steroids: β -sitosterol. Oxygen-containing heterocycles: irisoid A. Alkyl glycosides: ethyl- β - <i>D</i> -glucopyranoside [61].		<i>I. tenuifolia</i> is known as Gres-Ma in <i>Tibetan and Mongolian medicine</i> . The underground parts and seeds are used to treat intoxication, gastric colic, as an anthelmintic agent [53, 54]. The roots and rhizomes are commonly used in traditional Mongolian medicine for the treatment of kidney disorders. A root decoction is used for urethral stones, hypertension caused by adrenal gland diseases, and to relief renal colic and chronic nephritis [122]. Tenuifone, izalpinin and 5,7-dihydroxy-2';6-dimethoxyisoflavone have shown a considerable antioxidant activity [61]. Compounds 3'-hydroxy-5,7-dimethoxy-4-O-2'-cycloflavan and 5,2';3'-trihydroxy-6,7-dimethoxyflavanone have shown a neuroprotective activity [120]. Pretreatment with <i>I. tenuifolia</i> flavonoids from underground parts significantly inhibited H ₂ O ₂ -induced cell death in cortical neurons <i>in vitro</i> [123]. Treatment with ethanol extract of <i>I. tenuifolia</i> either 1 hr prior or immediately after middle cerebral artery occlusion (MCAO) significantly reduced the infarct volume in mice [124]. Flavonoids from rhizomes and roots have shown an antimicrobial activity against fungi, gram-positive and gram-negative bacteria. 5,7,2';3'-tetrahydroxyflavanone was found to possess the highest antibacterial activity against vancomycin resistant <i>Enterococcus faecalis</i> and <i>Mycobacterium vaccae</i> . Isolated compounds and plant crude extract <i>in vitro</i> showed potent inhibitory effect against leukemia cells. 5,2';3'-dihydroxy-6,7-dimethoxyisoflavone was active against both of the leukemia cells K-562 and THP-1 [121].	53, 54, 60, 61, 74, 120, 121, 122, 123, 124
Rhizomes Roots	Flavonoids and isoflavonoid: 5,2';3'-trihydroxy-6,7'-methylenedioxyflavanone, 5,2'-dihydroxy-6,7'-methylenedioxyflavanone, 5,2';3'-trihydroxy-7'-methoxyflavanone, 5,3'-dihydroxy-7,2'-dimethoxyflavanone [60], 3'-hydroxy-5,7'-dimethoxy-4-O-2'-cycloflavan, 3'-hydroxy-5-methoxy-6,7'-methylenedioxy-4-O-2'-cycloflavan, 5,2';3'-trihydroxy-6,7'-dimethoxyflavanone [120], 5-hydroxy-6,7'-dimethoxyisoflavone-2'-O- β - <i>D</i> -glucopyranoside, 5-methoxy-6,7'-methylenedioxy-4-O-2'-cycloflavan, (2S)-5,7,2';3'-tetrahydroxyflavanone, 5,2';3'-trihydroxy-6,7'-dimethoxyisoflavone, 5,7-dihydroxy-6,2'-dimethoxyisoflavone, 3,5,3'-trihydroxy-7,2'-dimethoxyflavanol [121]. Steroids: β -sitosterol. Macrolides: monitrstenulide [121].			
Aerial parts	Phenolic acids: vanillic, protocatechuic, <i>p</i> -coumaric, <i>m</i> -hydroxybenzoic, ferulic [74].			

Table 2. (Contd.)

[illegible]

Table 2. (Contd.)

Species	Plant part	Compounds	Biological activity and use	Ref.
<i>I. laevigata</i> Fisch. et C.A. Mey.	Rhizomes Roots	<i>Xanthones</i> : mangiferin [159].	<i>In Chinese</i> traditional medicine <i>rhizomes</i> are used as expectorant [113]. <i>In Transbaikalia</i> , root decoction is used for venereal diseases, as a laxative and emetic. <i>Flowers and fresh leaves</i> are used for snake bites (<i>Coluber natrix</i>) [40].	40, 56, 68, 69, 113, 144, 138, 159, 160, 161
	Aerial parts	<i>Xanthones</i> : mangiferin (0.73%) [69].		
	Leaves	<i>Phenolic acids from hydrolyzed</i> : <i>p</i> -coumaric, sinapic, ferulic, caffeic. <i>Anthocyanins</i> : delphinidin, cyanidin [68].		
	Flowers	<i>Anthocyanins</i> : malvidin-3-(<i>p</i> -coumaroyl)-rutinoside-5-glucoside (ensatin), petunidin-3-(<i>p</i> -coumaroyl)-rutinoside-5-glucoside (petanin), malvidin 3-rutinoside-5-glucoside, petunidin 3-rutinoside-5-glucoside, delphinidin-3-rutinoside-5-glucoside [56, 138]. <i>C-Glycosylflavone</i> : isovitexin, isovitexin 2"- <i>O</i> -rhamnoside, isovitexin 4'- <i>O</i> -glucoside, isovitexin 4'- <i>O</i> -galactoside, isovitexin X"- <i>O</i> -galactoside, acylated isovitexin X"- <i>O</i> -glucoside, vitexin, isoorientin, swertisin X"- <i>O</i> -rhamnoside, swertisin X"- <i>O</i> -galactoside, vicenin-2, isoscoparin, scoparin 7- <i>O</i> -glucoside, apigenin 6,8-di- <i>C</i> -glycoside [144, 160, 161].		
	Rhizomes Roots	<i>Carbohydrate</i> (%): sucrose (0.3), fructans (4.7) [55]. <i>Triterpenoids</i> : (6 <i>R</i> ,10 <i>S</i> ,11 <i>S</i> ,14 <i>S</i> ,26 <i>R</i> -(+)-29-Acetoxy-14,15-dihydro-26-hydroxyspiroirida-15(28),16-dienal; (6 <i>R</i> ,10 <i>S</i> ,11 <i>S</i> ,14 <i>S</i> ,26 <i>R</i> -(+)-14,15-Dihydro-26,29-dihydroxyspiroirida-15(28),16-dienal; (6 <i>R</i> ,10 <i>S</i> ,11 <i>R</i>)-26ξ-Hydroxy-13ξ-oxaspiroirid-16-enal; (6 <i>R</i> ,10 <i>S</i> ,11 <i>S</i>)-13ξ-oxaspiroirid-16-enal [162]; 26-acetoxy-22-methylene-irid-16-enal [162, 163], squalene, α-curcumene, sclareol, eugenol, α-bergamotene, [120]-farnesene, manoyloxide, <i>epi</i> -manoyloxide [164], γ-curcumene, α-bisabolol, β-bisabolene [165]. <i>Higher fatty acids</i> : palmitic, lauric, capric, caproic, myristic [165], ethyl linoleate, ethyl palmitate [164]. <i>Alkanes</i> : heptadecane, hexadecane [164]. <i>Organic acid</i> : shikimic, quinic, malic, succinic, citric, fumaric, lactic [55, 166, 167], gallic, 2,3,5-trihydroxybenzoic, gentisic, <i>p</i> -hydroxybenzoic [168], myristic [129, 169]. <i>Flavonoid</i> : 3,6-dimethoxy-5,7-dihydroxyflavone [168], kaempferol, quercetin [170], epigallocatechin, catechin, epicatechin, galocatechin [171], hyperoside [169]. <i>Flavones</i> : apigenin, formononetin, hispidulin [169, 170]. <i>Isoflavonoid</i> : ayamenin A, irilin B, iristectorigenin A, tectorigenin, 5,7-dihydroxy-2',6'-dimethoxyisoflavone, 5,7-dihydroxy-2',6'-dimethoxyisoflavone 7- <i>O</i> -β- <i>D</i> -glucopyranoside [168], 5,6-dihydroxy-7,8,3',5'-tetramethoxyisoflavone [63], irisolidone, irigenin, iridin [172], irilone, irisolone (nigricin) [173], iristectorigenin B (5,7,4'-trihydroxy-6,3'-dimethoxyisoflavone) [171], tectoridin, nigricin-4'-glucoside, 6,7-dihydroxyisoflavone, 5,7-dihydroxy-4'-methoxyisoflavone [169].		
<i>I. pseudacorus</i> L.	Rhizomes Roots		<i>The extracts of flowers, leaves, roots</i> are active against <i>Bact. abortus</i> Bang [199, 200]. <i>Essential oil</i> from <i>I. pseudacorus</i> contains antibacterial agents active against both gram-positive and gram-negative bacteria [201]. <i>Extracts of leaves and rhizomes</i> have been reported to have antibacterial activity against <i>Staphylococcus aureus</i> , <i>Escherichia coli</i> , <i>Pseudomonas aeruginosa</i> , <i>Bacillus subtilis</i> [202]. <i>Methanolic extracts from rhizomes</i> possessed antibacterial activity against human gram-positive <i>Staphylococcus aureus</i> and <i>Enterococcus faecalis</i> and gram-negative <i>Pseudomonas aeruginosa</i> and <i>Klebsiella pneumoniae</i> pathogens. <i>Extract from rhizome</i> effectively combats <i>Staphylococcus aureus</i> biofilm [171]. <i>I. pseudacorus</i> non-polar fractions showed potent biofilm inhibitory potential against biofilm formation of methicillin-resistant <i>Staphylococcus aureus</i> and methicillin susceptible <i>Staphylococcus aureus</i> [203]. <i>Methanol extract of I. pseudacorus</i> stimulated alkaline phosphatase activity and collagen type I secretion with a substantial effect on osteoblast proliferation [35]. <i>Rhizome extract</i> exhibits activity against human cancer cell lines, especially against estrogenpositive MCF-7 breast cancer cell line [171].	35, 36, 37, 38, 39, 41, 51, 55, 63, 67, 68, 71, 78, 114, 125, 126, 129, 137, 162, 163, 164, 165, 166, 167,

Table 2. (Contd.)

Species	Plant part	Compounds	Biological activity and use	Ref.
		Xanthones: mangiferin (2-(C-β-D-glucopyranosyl)-1,3,6,7-tetrahydroxyxanthone), isomangiferin (4-(C-β-D-glucopyranosyl)-1,3,6,7-tetrahydroxyxanthone), norathyrin (1,3,6,7-tetrahydroxyxanthone) [174], daidzein, genistein, genistein-7-glucoside [169–171]. Amino acid: methyl-2-pyrrolidone-5-carboxylate, 2-pyrrolidone-5-carboxylic [175], 3-(3-carboxyphenyl)alanine [176, 177], aspartic, lysine, glutamic, arginine, glycine, leucine, threonine, valine, phenylalanine, serine, alanine, isoleucine, histidine, tyrosine, methionine, proline [178]. Hydroxybenzoic acid: gallic, 2,3,5-trihydroxybenzoic, gentisic, <i>p</i>-hydroxybenzoic, protocatechuic [168, 171, 179]. Phenolcarboxylic acids: caffeic, ferulic, chlorogenic, neochlorogenic [170], <i>p</i>-coumaric, <i>trans</i>-cinnamic [169], ellagic [171]. Polysaccharides: water-soluble polysaccharides, pectic substances, hemicellulose A, hemicellulose B. Monosaccharides: glucose, xylose, galactose, mannose, rhamnose, arabinose [39]. Coumarins: coumarin, umbelliferone (7-hydroxycoumarin), herniarin (7-methoxycoumarin), daphnoretin, esculetin, scopoletin, isoscapoletin [180].	In India, <i>I. pseudacorus</i> is used in dysmenorrhoea and leucorrhoea. Juice of the root is used for obstinate coughs and convulsions [129]. In traditional medicine of Belarus, a decoction of the rhizomes is used for gastric ulcers, as a hair growth stimulant [204]. In Ukrainian traditional medicine, peeled and cut rhizomes are used as an emetic, laxative, diuretic, stimulant, expectorant and coating agents. Powder from dried rhizomes is used as a toothpaste ingredient, to relieve toothache in children [37, 41, 174]. In Bulgaria, the rhizomes were used for diabetes [205]. In Russian traditional medicine, a decoction of the rhizomes was used for respiratory infections, pneumonia, diseases of the urinary organs, ascites, scurvy, scrofula, dropsy and epilepsy, as an analgesic for headache and toothache, obstetric, hemostatic for algomenorrhea and metrorrhagia; as a laxative and antidiarrhoeal agent [36, 38, 206, 207]. Rhizomes were used for snake bites [41]. The sap from the rhizomes was used (orally) for epilepsy; locally – for diabetes, toothache, stomatitis [51, 206]. Seeds were used as a coffee substitute [51]. Rhizomes is one of the ingredients of M.N. Zdrenko polyherbal tea used in anticancer therapy [208, 209]. Rhizomes were included in different pharmacopoeias and have been used as a strong purgative agent [210]. In homeopathic medicine, extracts of rhizomes are used for different forms of psychic ailments, migraine, for neuralgia of the trigeminus, in hyperemesis and hyperacidity of the stomach [163, 211]. <i>I. pseudacorus rhizome sap</i> was used by English rural population and administered with buckthorn syrup every one-two hours to treat dropsy [212]. <i>I. pseudacorus</i> was used externally for toothache, scabs and scrofula, internally for hydrops [78]. Dry hydrophilic leaf extract showed pronounced diuretic activity [213]. Underground parts and flowers were used as a tanning agent and yellow dye for leather, wool, paper [126, 214–217]. The larvicidal activity of ethanol leaf extracts on the larvae of <i>Culex pipiens</i> and <i>Aedes caspius</i> mosquitoes were investigated. The miracicidal and cercaricidal properties were tested against <i>Schistosoma mansoni</i> miracidia [218]. Floating treatment wetlands (FTWs) planted with <i>I. pseudacorus</i> uptake nitrogen and phosphorous, and can be used in a temperate climate to overcome problems with excessive algae growth in surface waters in urban and agricultural settings [219].	168, 169, 170, 171, 172, 173, 174, 175, 176, 177, 178, 179, 180, 181, 182, 183, 184, 185, 186, 187, 188, 189, 190, 191, 192, 193, 194, 195, 196, 197, 198, 199, 200, 201, 202, 203, 204, 205, 206, 207, 208, 209, 210, 211, 212,
	Leaves	Isoflavonoid: ayamenin A, ayamenin B, iriline B, irilin C, biochanin A, tectorigenin, irilin A, iristectorigenins A, iristectorigenins B, 3'-<i>O</i>-methylorobol, pratensein, genistein [181], ayamenins C, ayamenins D, ayamenin E [182], 5,7,3'-trihydroxy-6-methoxycoumaronochromone, lupinalbin A [181]. Flavanol: galloctechin-4-β-ol [67, 71]. Flavanones: 5,7,2'-trihydroxyflavanone [182]. Dihydroflavonol: alpinone [182], 7-<i>O</i>-methyl(dihydrokaempferol (aromadendrin 7-methyl ether). Flavones: apigenin, luteolin, hispidulin [182, 183]. Anthocyanins: delphinidin, cyanidin [68]. Catechins: (-)-epicatechin. C-glycosides: isovitexin, vitexin, isoorientin, orientin, vicenin-II. Organic acid: protocatechuic, gallic [183]. Xanthones: mangiferin, isomangiferin, mangiferin <i>O</i>-glycoside [71, 183, 184]. Higher fatty acids: myristic, stearic, palmitic, lignocenic, arachidic, behenic, myristoleic, oleic, linoleic, linolenic, erucic, palmitoleic [185]. Phenolic acids: <i>p</i>-coumaric, sinapic, ferulic [68]. Ascorbic acid 80–140 mg% [125].		
	Aerial parts	Isoflavonoid: 5,7-dihydroxy-2',6-dimethoxyisoflavone, irilone [186]. Flavanol: 3,6-dimethoxy-5,7-dihydroxyflavone [186]. Dihydroflavonol: alpinone, <i>trans</i>-3-hydroxy-5,7-dimethoxyflavanone [186].		

Table 2. (Contd.)

Species	Plant part	Compounds	Biological activity and use	Ref.
<i>I. sanguinea</i> Donn. (= <i>I. orientalis</i> Thunb., <i>I. nerzhinskia</i> Lodd. f. <i>albiflora</i> Honda.	Flowers	Flavonol : isorhamnetin, kaempferol [187]. Carotenoids : β -carotene, lutein, zeaxanthin, violaxanthin [137, 188, 189]. Tannins : 2.82% pyrogallol derivatives [114, 190]. Higher fatty acids : stearic, palmitic, oleic, linoleic, linolenic [191]. The content of lipids has been investigated [191].		213, 214, 215, 216, 217, 218, 219
	Seeds	Quinones : irisquinone; 2-methoxy-6-(pentadec-8-enyl)-1,4-benzoquinone; 2-methoxy-6-pentadecyl-1,4-benzoquinone; 2-methoxy-6-nonadecyl-1,4-benzoquinone; 6-(heptadec-10'-enyl)-5-hydroxy-2-methoxy-1,4-benzoquinone; 5-hydroxy-2-methoxy-6-(pentadecenyl)-1,4-benzoquinone; 5-hydroxy-2-methoxy-6-pentadecyl-1,4-benzoquinone; 3-methoxy-5-(heptadec-10'-enyl)phenol; 3-methoxy-5-(pentadecenyl)phenol; 3-methoxy-5-pentadecylphenol; 5-(heptadec-10'-enyl)-3,4-dimethoxyphenol; 3,4-dimethoxy-5-(pentadecenyl)phenol; 3,4-dimethoxy-5-pentadecylphenol; 3,4-dimethoxy-5-(nonadecenyl)phenol [192–195]. Carbohydrates : glucose, fructose, sucrose, raffinose, stachyose [196]. The content of lipids has been investigated [197].		
	Callus culture	Phenylpropanoids : lavandoside. Neolignan : dehydrodiconiferyl alcohol-4- <i>O</i> - β - <i>D</i> -glucopyranoside. Isoflavonoids : tectoridin, tectorigenin, iristectorigenin A [198].		
<i>I. sanguinea</i> Donn. (= <i>I. orientalis</i> Thunb., <i>I. nerzhinskia</i> Lodd. f. <i>albiflora</i> Honda.	Rhizomes	Carbohydrate (%): sucrose (1.6), fructans (7.0), starch (3.2) [55].	<i>In Chinese</i> traditional medicine rhizomes and roots are used to treat furuncles, carbuncles, stomach ache, indigestion, bruises, hemorrhoids, scabies, edema, cirrhotic ascites, difficult urination, constipation [148]. Flowers, seeds and rhizomes have been used in <i>Tibetan medicine</i> [42]. The <i>Nunais</i> used decoction of rhizomes for gynecological diseases and as an emetic and laxative [43]. <i>In Russian</i> traditional medicine rhizomes were used for dropsy and as an anthelmintic agent, externally – as a remedy for burns [24].	24, 42, 43, 55, 56, 71, 95, 148, 220, 221, 222, 223
	Roots			
	Leaves	Flavonols : myricetin, quercetin [71], myricetin 3- <i>O</i> -rhamnoside, quercetin 3- <i>O</i> -rhamnoside [220]. C-Glycosylflavones : swertiajaponin, swertisin 2"- <i>O</i> -glucoside (flavayamenin), swertisin, isoorientin 2"- <i>O</i> -glucoside, acetylated swertisin 2"- <i>O</i> -glucosides, acetylated swertisin [220].		
<i>I. sanguinea</i> Donn. (= <i>I. orientalis</i> Thunb., <i>I. nerzhinskia</i> Lodd. f. <i>albiflora</i> Honda.	Flowers	C-Glycosylflavones : swertiajaponin, luteoyamenin, swertisin, swertisin 2"- <i>O</i> -glucoside (flavayamenin) [95, 221, 222], embigenin 2"- <i>O</i> -rhamnoside [223], vicienin-2, swertiajaponin X"- <i>O</i> -hexoside, isoorientin, isovitexin 2"- <i>O</i> -glucoside, isovitexin [220]. Anthocyanins : delphinidin-3-(<i>n</i> -coumaroyl)-rutinoside 5-glucoside, petunidin-3-(<i>p</i> -coumaroyl)-rutinoside-5-glucoside (petanin), delphinidin-3-rutinoside-5-glucoside, petunidin 3-rutinoside-5-glucoside, malvidin glycoside [56], delphinidin 3- <i>O</i> -(4"- <i>p</i> -coumaroylrhamnosyl)-(1→6)-glucoside]-5- <i>O</i> -glucoside, petunidin 3- <i>O</i> -(4"- <i>p</i> -coumaroylrhamnosyl)-(1→6)-glucoside]-5- <i>O</i> -glucoside [220].		

Table 2. (Contd.)

Species	Plant part	Compounds	Biological activity and use	Ref.
<i>I. sibirica</i> L.	Rhizomes	Carbohydrate (%): sucrose (2.3), fructans (2.7), starch (2.5) [55]. Xanthones : mangiferin, daidzein, genistein-7-glucoside [50, 169], formononetin, isomangiferin [170]. Phenolcarboxylic acids : caffeic, gallic, chlorogenic, <i>p</i> -coumaric, ferulic, <i>trans</i> -cinnamic [50, 170].	<i>I. sibirica</i> is considered as an antisyphilitic treatment [78]. <i>In Tibetan</i> medicine, underground parts, flowers and seeds were used for pneumonia, bronchitis, gastritis, gynecological diseases [20]. <i>In Russian</i> traditional medicine underground parts were used for ascites, syphilis, scurvy, as a hemostatic, laxative, anthelmintic [146, 230]. <i>In Siberia</i> , a decoction of the herb was used for heart disease, during pregnancy, for gastrointestinal diseases, and for anthrax [231]. The fruits were used as decoction, and as a wound healing remedy [41]. <i>In the Urals</i> , a whole plant with fruits as decoction or alcoholic infusion was used orally for gastrointestinal diseases and dysentery. The decoction was taken by women after childbirth, and externally as a wound healing agent [232]. <i>In Transbaikalia</i> root decoction was used to treat venereal diseases, as a laxative and emetic agent. Flowers and fresh leaves were used for snake bites (<i>Coluber natrix</i>) [40]. The flowers were used as a green dye [233]. Flower pigments were used as a natural textile dye [234]. <i>In Brazil</i> <i>I. sibirica</i> is known as “Palmeirinha” and its roots are used for diarrhea [235]. Dry and lipophilic extracts from the leaves and rhizomes exhibited an antibacterial activity against <i>Staphylococcus aureus</i> , <i>Escherichia coli</i> , <i>Proteus vulgaris</i> , <i>Pseudomonas aeruginosa</i> , <i>Bacillus subtilis</i> and <i>Candida albicans</i> [236].	6, 20, 40, 41, 50, 55, 68, 71, 78, 125, 146, 169, 170, 224, 225, 226, 227, 228, 229, 230, 231, 232, 233, 234, 235, 236
	Roots	Flavonoids and isoflavonoid : germanaism B, tectoridin, irisolidone- <i>D</i> -glucoside, nigricin, irigenin, nigricin-4'-glucoside, 6,7-dihydroxyisoflavone, iristectorigenin B, nigricin, 7-hydroxyisoflavone, 5,6-dihydroxy-7,8,3',5'-tetramethoxyisoflavone, 5,7-dihydroxy-4'-methoxyisoflavone [50, 169]. Flavonols : hyperoside, isoquercetin, kaempferol [169], quercetin [170]. Flavones : apigenin. C-Glycosylflavones : apigenin-7-glucoside [169]. Triterpenoids : (6 <i>R</i> ,10 <i>S</i> ,11 <i>S</i>)-(+)-26-hydroxyiridal, (6 <i>R</i> ,10 <i>S</i> ,11 <i>S</i>)-(+)-17 ξ ,26-dihydroxyiridal, (6 <i>R</i> ,10 <i>S</i> ,11 <i>S</i>)-(+)-17 ξ -hydroxyiridal, (6 <i>R</i> ,10 <i>S</i> ,11 <i>S</i>)-(+)-10-deoxy-17 ξ -hydroxyiridal [224], 10-deoxy-17-hydroxyiridal, (6 <i>R</i> ,10 <i>S</i> ,11 <i>S</i>)-17,29-didehydroiridal [225]. Higher fatty acids : lauric, myristic, palmitic, palmitoleic, stearic, oleic, linoleic, 9-methyl tetradecenoic, isomyristic, pentadecanoic, 14-methyl palmitoleic, heptadecanoic, 10-octadecadienoic, linolenic, arachic, behenic, tricosyl, lignoceric [6].		
	Flowers	C-Glycosylflavones : 6- <i>C</i> -glucosyl-7- <i>O</i> -methylapigenin (swertisin), 6- <i>C</i> -glucosyl-7- <i>O</i> -methyluteolin (swertiajaponin), 6- <i>C</i> -sophorosyl-7- <i>O</i> -methylapigenin (flavoayamenin, spinosin), 6- <i>C</i> -sophorosyl-7,4'-di- <i>O</i> -methylapigenin (embinoidin) [226].		
	Leaves	Xanthones : mangiferin [227]. Flavonols : quercetin, myricetin [71, 228], epigallocatechin, catechin [227]. Anthocyanins : delphinidin, cyanidin. Phenolic acids : caffeic, <i>p</i> -coumaric, sinapic, ferulic [20, 68, 227]. Higher fatty acids : lauric, palmitic, palmitoleic, stearic, oleic, linoleic, pentadecanoic, heptadecanoic, linolenic, arachic, behenic, tricosyl, lignoceric, hexadecadienoic [6]. Isoflavones : tectorigenin-7- <i>O</i> -hexosyl-4'- <i>O</i> -hexoside, tectorigenin-7- <i>O</i> -dihexoside, dichotomitin, genistein- <i>O</i> -hexoside (genistin), genistin <i>O</i> -hexoside isomer, genistein, tectorigenin, rhamnocitrin, isotectorigenin, tectorigenin- <i>O</i> -hexoside (tectoridin), iristectorigenin B- <i>O</i> -hexoside (iristectorin B), iristectorigenin A- <i>O</i> -hexoside (iristectorin A), iristectorigenin A, iristectorigenin B, irisolidone, irisolidone isomer, irisolidone <i>O</i> -hexoside, irisolidone <i>O</i> -hexoside isomer, irilone, iristectorin B-4'- <i>O</i> -glucoside, irigenin [227]. Flavones : naringenin-7- <i>O</i> -dihexoside (naringin), naringenin, eupatilin, hispidulin, genkwanin (7-methoxyapigenin), dihydrokaempferol, 6,7,3',4'-tetrahydroxyflavone [227]. C-Glycosylflavones : apigenin- <i>C</i> -hexoside (vitexin), 4'- <i>O</i> -methyl-apigenin 8- <i>C</i> -hexoside, apigenin 8- <i>C</i> -(2"-hexosyl)-hexoside [227]. O-glycosylflavones : flavonoid <i>O</i> -dihexoside, chrysoeriol-7- <i>O</i> -hexoside [227]. Ascorbic acid 400 mg% [125].		
	Seeds	Carbohydrates : polysaccharides were composed of D-glucose, D-mannose, D-galactose [229]. Polysaccharide : glucomannan (18%) [55].		

Table 2. (Contd.)

Species	Plant part	Compounds	Biological activity and use	Ref.
Triptelatae Alexeeva				
I. setosa Pall.	Ariel parts	Xanthones: mangiferin (2.76%) [69].	In Russian traditional medicine a decoction of the rhizomes was used for pneumonia, ascites, and sore throat; externally – for wounds, ulcers, fistulas, and to remove freckles [46]. In Yakutia, the rhizomes were used as an analgesic (for toothaches) and emetic agents. Dried rhizomes were used as an insecticide. A decoction of the rhizomes was used for gastrointestinal diseases [240]. In Japanese medicine, the powdered rhizomes were used for scabies [45]. Seeds were used as a coffee substitute [46]. The rhizomes were used as a yellow dye [46].	45, 46, 56, 69, 71, 144, 161, 237, 238, 239, 240
	Leaves	C-Glycosylflavones: luteolin 6,8-di-C-glycoside, swertiajaponin, isoswertiajaponin X"-O-rhamnoside, isoswertisin, swertisin X"-O-rhamnoside, isoorientin, isovitexin, orientin, vitexin, vicenin-2, isovitexin 4'-O-glucoside, swertisin [237, 238]. Xanthones: mangiferin, isomangiferin [71, 237].		
	Flowers	Anthocyanin: malvidin 3-O-[(4"-p-coumaroylrhamnosyl)-(1→6)-glucoside]-5-O-glucoside, malvidin glycoside, petunidin 3-O-[(4"-p-coumaroylrhamnosyl)-(1→6)-glucoside]-5-O-glucoside, delphinidin hexoside [237, 239], delphinidin 3-O-rutinoside-5-O-glucoside, delphinidin 3-O-(p-coumaroylrutinoside)-5-O-glucoside, petunidin 3-O-rutinoside-5-O-glucoside, malvidin 3-O-rutinoside-5-O-glucoside [56, 161]. C-Glycosylflavones: isovitexin 4'-O-galactoside, vitexin 4'-O-galactoside, isovitexin X"-O-glucoside, swertisin 2"-O-glucoside, isoorientin, isovitexin, orientin, vitexin, vicenin-2, isovitexin 4'-O-glucoside, swertisin [144, 237].		
Ioniris (Spach) Rodion				
I. ruthenica Ker-Gawl.	Rhizomes	Carbohydrate (%): sucrose (1.0), fructans (1.2), starch (3.3) [55]. Content of polysaccharides 1.33% and pectin 2.2% [241]. The presence of saponins [242] and flavonoids [243] has been investigated.	Roots, flowers and seeds are used in Tibetan medicine [146]. The flowers are used as an anthelmintic agent [47]. A decoction of the roots, pre-dried, are used for gynecological diseases [244].	47, 55, 68, 146, 241, 242, 243, 244
	Leaves	Phenolic acids from hydrolyzed: p-coumaric, sinapic, ferulic [68]. Content of polysaccharides 3.5% and pectin 3.1% [241].		
Oncocylus (Siemss.) Baker				
I. acutiloba C. A. Mey (I. ewbankiana Foster)	Rhizomes	Carbohydrate (%): sucrose (2.5), starch (11.8) [55]. Higher fatty acids: capric, lauric, myristic, stearic, palmitic, oleic, linoleic, linolenic [245].	The essential extract exhibits antibacterial activity [246].	55, 125, 245, 246
	Leaves	Ascorbic acid 280–391 mg% [125].		
Iris				
I. aphylla L. (= I. hungarica Waldst. et Kit.)	Rhizomes	Xanthones: mangiferin, isomangiferin [50, 170, 247]. Phenolcarboxylic acids: n-coumaric, chlorogenic, cinnamic, caffeic [50], trans-cinnamic [169], gallic [170]. Flavonoids and isoflavonoid: 7,4'-dihydroxyisoflavone (daidzein), 7-hydroxy-4'-methoxyisoflavone (formononetin), 5,7,4'-trihydroxyisoflavone (genistein), 5,7,4'-trihydroxy-6-methoxyisoflavone (tectorigenin), 4',5'-dihydro-6-methoxy-7-(O-β-D-glucoside)isoflavone (tectordin), germanaism B, irisolidone-D-glucoside, irigidone, nigricin, irigenin, 5,6-dihydroxy-7,8,3',5'-tetramethoxyisoflavone [50, 247], 5,7,4'-trihydroxy-6,3'-di-methoxyisoflavone (iristectorigenin B), iristectorigenin B-7-O-β-D-glucoside (iristectorin B) [170], nigricin-4'-glucoside [169, 170].	The essential extract exhibits antibacterial activity [48, 49]. 1% dry rhizome extract has shown the highest inhibitory activity for gram-positive bacteria and fungi [247]. The dry extract of the leaves at the concentration 1% has a more pronounced antibacterial activity than the extract of the rhizomes and is the most susceptible to gram-negative bacteria – Escherichia coli, Proteus vulgaris, Pseudomonas aeruginosa. The lipophilic extracts of the leaves and rhizomes were more susceptible to gram-positive bacteria [236].	48, 49, 50, 68, 125, 169, 170, 236, 247, 248, 249
	Roots			

Table 2. (Contd.)

Species	Plant part	Compounds	Biological activity and use	Ref.
	Leaves	Xanthone: mangiferin [68]. Phenolic acids: <i>p</i>-coumaric, ferulic [68], syringic, <i>p</i>-hydroxybenzoic, vanillic [248]. Carboxylic acids: oxalic, malonic, fumaric, methoxysuccinic, succinic, 3-oxo-2-methylglutaric, malic, suberinic, azelaic, citric [248]. Ascorbic acid 450 mg% [125].	Aqueous and ethanol extract of rhizomes showed a potent antioxidant capacity, anti-inflammatory and antiallergic activity. Aqueous extract of rhizomes showed inhibitory effect on the lipid droplets in Huh7 liver cells. Ethanol extract of rhizomes was effective against both melanoma (ICR39) and triple-negative breast cancer (MDA-MB-231) cells [50]. Dry extracts of leaves and rhizomes showed an anabolic activity, reduced urea excretion, normalized metabolism, restored nitrogen balance, and inhibited protein catabolism in rats [249].	41, 51, 52, 55, 69, 71, 125, 250, 251
<i>I. pumila</i> L.	Rhizomes	Carbohydrate (%): sucrose (2.0), starch (5.0) [55]. Xanthones: iriflophenone 4-<i>O</i>-hexoside, 7-<i>O</i>-methyl-mangiferin, isomangiferin, 7-<i>O</i>-methyl-isomangiferin, iriflophenone, 4-<i>O</i>-methyl-iriflophenone, bellidifolin [250]. O-glycosylflavones: luteolin 7-<i>O</i>-(2"-<i>p</i>-coumaroyl)-rhamnoside.	In Russian traditional medicine the rhizomes were used as antifever, emetic [41], and antirabies agent [51]. Rhizomes and leaves show antifungal activity [52].	
	Roots	Isoflavones and derivatives: tectoridin, iristectorin B, iristectorin A, iridin, 7-<i>O</i>-methyl-tectorigenin 4'-<i>O</i>-(6"-hexosyl)-hexoside, irifloside, irilone 4'-<i>O</i>-hexoside, irisolidone 7-<i>O</i>-hexoside, tectorigenin, iristectorigenin A, irigenin, irilone, iriflogenin, irisolidone [250]. Content (mg/g of dry weight) of soluble sugars 15.5, starch 6.3 [250].		
	Aerial parts	Xanthones: mangiferin (1.28%) [69, 71], isomangiferin, 7-<i>O</i>-methyl-isomangiferin, iriflophenone, 4-<i>O</i>-methyl-iriflophenone, bellidifolin [250]. C-Glycosylflavones: luteolin 6-<i>C</i>-glucoside, apigenin 8-<i>C</i>-glucoside, apigenin 6-<i>C</i>-hexoside, 4'-<i>O</i>-methyl-apigenin 8-<i>C</i>-hexoside, 4'-<i>O</i>-methyl-apigenin 6-<i>C</i>-hexoside [250]. O-glycosylflavones: luteolin 7-<i>O</i>-(2"-<i>p</i>-coumaroyl)-rhamnoside [250]. Isoflavones and derivatives: iristectorin B, iristectorin A, tectorigenin, irilone 4'-<i>O</i>-(3-hydroxy-3-methylglutaryl)-hexoside, iristectorigenin A, irigenin [250]. Content (mg/g of dry weight) of soluble sugars 18.9, starch 2.3 [250]. Ascorbic acid 411–808 mg% [125].		
	Flowers	Xanthones: mangiferin, isomangiferin, 7-<i>O</i>-methyl-isomangiferin, iriflophenone [250]. C-glycosylflavones: luteolin 8-<i>C</i>-hexoside, luteolin 6-<i>C</i>-glucoside, apigenin 8-<i>C</i>-(2"-hexosyl)-hexoside, apigenin 8-<i>C</i>-(2"-pentosyl)-hexoside, apigenin 8-<i>C</i>-glucoside, apigenin 6-<i>C</i>-hexoside [250]. O-glycosylflavones: kaempferol 7-<i>O</i>-(6"-rhamnosyl)-hexoside [250]. Isoflavones and derivatives: irisolidone 7-<i>O</i>-hexoside [250]. Content (mg/g of dry weight) of soluble sugars 22.6, starch 1.17 [250]. The composition and relative contents of floral scent components has been investigated [251].		

Table 2. (Contd.)

Species	Plant part	Compounds	Biological activity and use	Ref.
<i>I. potaninii</i> Mazim.	Under-ground parts	<i>Psammiris</i> (Spach) Taylor Isoflavone: 7,8,5'-trihydroxy-6,3',4'-trimethoxyisoflavone; 7,4'-dimethoxy-8,3',5'-trihydroxy-6- <i>O</i> - β - <i>D</i> -glucopyranosylisoflavone; 3',4',5'-tetramethoxy-6,7-methylenedioxyisoflavone (iris-florentin); 3'-methoxy-4',5'-dihydroxy-6,7-methylenedioxyisoflavone (iriskumaonin); 3',4'-dimethoxy-5',5'-dihydroxy-6,7-methylenedioxyisoflavone (dichotomitin); 4',5'-dimethoxy-3-hydroxy-6,7-methylenedioxyisoflavone (iriflogenin); 5'-methoxy-4'-hydroxy-6,7-methylenedioxyisoflavone irisolone (nigricin) [4, 5, 195, 252, 253]. Flavanol: 5,3',3'-trihydroxy-7,4'-dimethoxyflavanone (blumeatin B) [4]. Xanthones: iriflophenone [4, 254].	<i>In Mongolian</i> traditional medicine roots are used for worm and poisoning diseases, wound healing, eye yellowing, drying lymph leakage, and treats stomach and large intestine aches, bacterial infections, cancer, and inflammation. Extract of the underground parts exhibit nephroprotective activity [4, 254, 255].	4, 5, 195, 252, 253, 254, 255
	Rhizomes	Carbohydrate (%): sucrose (2.3), starch (4.1) [55]. Xanthones: isomangiferin, iriflophenone 4- <i>O</i> -(6"-acetyl)-hexoside, polygalaxanthone III, 7- <i>O</i> -methyl-isomangiferin, iriflophenone, 4- <i>O</i> -methyl-iriflophenone [250]. O-glycosylflavones: irisdichotin B [250]. Isoflavones and derivatives: irilone 4'- <i>O</i> -(6"-hexosyl)-hexoside, iridin, dichotomitin 3'- <i>O</i> -(6"-hexosyl)-hexoside, 7- <i>O</i> -methyl-tectorigenin 4'- <i>O</i> -(6"-hexosyl)-hexoside, nigracin 4'- <i>O</i> -(6"-hydroxy-3-methylglutaryl)-hexoside, irifloside, irilone 4'- <i>O</i> -hexoside, irisolidone 7- <i>O</i> -hexoside, tectorigenin, dichotomitin 3'- <i>O</i> -hexoside, irigenin [250]. Phenolic acids: protocatechuic, <i>p</i> -hydroxybenzoic, vanillic, syringic, <i>p</i> -coumaric, ferulic [74]. Content (mg/g of dry weight) of soluble sugars 9.9, starch 12.0 [250].	The infusion was used as lotion for breast tumors [256]. <i>In Tibetan</i> medicine, the underground parts and seeds are used for intoxication, gastric colic, as an anthelmintic and antibacterial agents [53, 54]. <i>In Mongolian</i> medicine, flowers and roots were used as a hemostatic agent (externally), for blood diseases, and osteoalgia [27, 257]. Rhizomes were used for sepsis, various infectious diseases and as an external hemostatic agent [9, 258].	9, 27, 53, 54, 55, 68, 71, 74, 250, 256, 257, 258
<i>I. humilis</i> Georgii (= <i>I. flavissima</i> Pull.)	Aerial parts	Xanthones: isomangiferin [250]. O-glycosylflavones: kaempferol 3- <i>O</i> -galactoside, kaempferol 3- <i>O</i> -glucoside [250]. Isoflavones and derivatives: iristectorigenin A 7- <i>O</i> -hexuronide, tectorigenin, iristectorigenin A [250]. Content (mg/g of dry weight) of soluble sugars 29.7, starch 1.1 [250].		
	Leaves	Flavanol: quercetin [68]. Xanthone: mangiferin [71]. Anthocyanins: cyanidin [68]. Phenolic acids: caffeic, <i>p</i> -coumaric, sinapic, ferulic [68].		
	Flowers	Xanthones: mangiferin, isomangiferin [250]. O-glycosylflavones: quercetin 3- <i>O</i> -galactoside, isorhamnetin 3- <i>O</i> -(2"-rhamnosyl)-hexoside, quercetin 3- <i>O</i> -glucoside, isorhamnetin 3- <i>O</i> -(6"-rhamnosyl)-hexoside, kaempferol 3- <i>O</i> -galactoside, kaempferol 7- <i>O</i> -(6"-rhamnosyl)-hexoside, isorhamnetin 3- <i>O</i> -galactoside, isorhamnetin 3- <i>O</i> -glucoside [250]. Content (mg/g of dry weight) of soluble sugars 23.0, starch 1.1 [250].		

In more detail, the pharmacological and useful properties of *Iris* species are presented in Table 2. Moreover, it contains data on the compound activity and occurrence in different plant part. A varied content of secondary metabolites in irises increases their medicinal importance.

Almost all flavonoid derivatives are present in all plant parts: in rhizomes, leaves, flowers and seeds of *I. lactea*, *I. oxipetala*, *I. tenuifolia*, *I. pseudacorus*, *I. sanguinea*, *I. sibirica*, *I. setosa*, *I. aphylla*, *I. pumila*, *I. potaninii*, *I. humilis* [15, 16, 35, 55–67].

It is worth to mention *I. aphylla*, *I. lactea* s.l., *I. pseudacorus*, *I. ruthenica*, *I. setosa*, *I. ensata*, *I. laevigata* from the aboveground parts of which mangiferin was isolated [13, 68–71].

Species of the genus *Iris* biosynthesize secondary metabolites of various classes, of which isoflavonoids and flavonoids are especially important. Such polyphenolic compounds most often act as phytoalexins, protecting plants from adverse effects of the environment [72].

Thus, this review shows that the genus *Iris* belongs to the number of genera are actively studied in the context of the chemical diversity and biological acti-

vity. It should be recognized that most of the information is related to *I. lactea* s.l., *I. oxypetala*, *I. ensata*, *I. pseudacorus* and *I. tenuifolia* and, to a lesser extent, to other species of the genus. Nevertheless, the wide spectrum of biological activity in some *Iris* species in presence of a very diverse set of biologically active compounds makes this genus promising for further research. It can be assumed that closely related species, which are poorly or not studied, have similar properties: *I. colchica* Kem.-Nat., *I. furcata* Bieb., *I. glaucescens* Bunge, *I. klattii* Kem.-Nat., *I. lokiae* Alexeeva, *I. loczyi* Kanitz, *I. uniflora* Pall.

Therefore, based on the analysis of the literature data, a wide range of pharmacological effects of the species of the genus *Iris* in Russia has been established. Species promising for further in-depth chemical and pharmacological study have been identified.

ACKNOWLEDGEMENTS

The present study was carried out within the framework of the Komarov Botanical Institute of the Russian Academy of Sciences research project № 122011900031-0.

REFERENCES

1. Mathew B. 1981. The *Iris*. London. 202 p.
2. Rodionenko G.I. 1987. The genus *Iris* L. (questions of morphology biology, evolution and systematics). London. 222 p.
3. Alexeeva N.B. 2020. *Iris* L. (Iridaceae Juss.) in Russia. Saint-Petersburg. 232 p. (In Russian, English)
4. Purev O., Purevsuren C., Narantuya S., Lkhagvasuren S., Mizukami H., Nagatsu A. 2002. New isoflavones and flavanol from *Iris potaninii*. — Chem. Pharm. Bull. (Tokyo). 50(10): 1367–1369. <https://doi.org/10.1248/cpb.50.1367>
5. Kaššák P. 2012. Secondary metabolites of the chosen genus *Iris* species. — Acta Univ. Agric. Silv. Mendelianae Brun. 60(8): 32. <https://doi.org/10.11118/actaun201260080269>
6. Mykhailenko O., Kovalyov V., Kovalyov S., Toryanik E., Osolodchenko T., Buidin Y. 2017a (2018). Fatty acid composition of lipids of *Iris sibirica*. — Čes. slov. Farm. 66(5): 220–226. <https://www.prolekare.cz/en/journals/czech-and-slovak-pharmacy/2017-5-7/fatty-acid-composition-of-lipids-of-iris-sibirica-63385>
7. Thunberg C.P. 1794. Botanical observation on the flora Japonica. — Trans. Linn. Soc. Lond. 2: 326–342. <https://doi.org/10.1111/j.1096-3642.1794.tb00268.x>
8. Grubov V.I. 1970. [Critical Remarks on the taxonomy and nomenclature of some species of iris. Flora of the USSR]. — Novitates systematicae plantarum vascularium. 6: 29–37. (In Russian)
9. Schroeter A.I. 1975. [Medicinal flora of the Soviet Far East]. Moscow. 328 p. (In Russian) <https://www.booksite.ru/fulltext/rusles/lekflora/index.htm>
10. Bunge A.A. 1833. Enumeratio plantarum, quas in China boreali collegit Dr. Al. Bunge: Anno 1831. St. Petersburg. 79 p.
11. Blinova K.F., Stukkey K.L. 1960. [Pharmacognostic study of the medicinal plants in Tibetan medicine: I]. — In: [Trudy Leningradskogo Khimiko-Farmatsevticheskogo Instituta. Voprosy Farmakognozii. 1]. Vol. 12. Leningrad. P. 135–155. (In Russian)
12. Blinova K.F., Stukkey K.L. 1964. [Pharmacognostic study of the medicinal plants in Tibetan medicine: II]. — In: [Trudy Leningradskogo Khimiko-Farmatsevticheskogo Instituta. Voprosy Farmakognozii. 2]. Vol. 17. Leningrad. 173–190. (In Russian)
13. Blinova K.F., Kalyupanova N.I. 1974. Xanthone glycosides of *Iris ensata*. — Chem. Nat. Compd. 10(4): 551. <https://doi.org/10.1007/BF00563849>
14. Blinova K.F., Glyzin V.I., Pryakhina N.I. 1977. A C glycoside from *Iris ensata*. — Chem. Nat. Compd. 13(1): 104. <https://doi.org/10.1007/BF00566189>
15. Pryakhina N.I., Blinova K.F. 1979. Phenolic acids of *Iris ensata*. — Chem. Nat. Compd. 15: 768. <https://doi.org/10.1007/BF00565598>

16. Pryakhina N.I., Blinova K.F. 1984. Luteolin C-glycosides from *Iris ensata*. — Chem. Nat. Compd. 20(1): 107. <https://doi.org/10.1007/BF00574811>
17. Pryakhina N.I., Sheichenko V.I., Blinova K.F. 1984. Acylated C-glycosides of *Iris lactea*. — Chem. Nat. Compd. 20(5): 554–559. <https://doi.org/10.1007/BF00580064>
18. [Plant Resources of Russia and neighboring states: Flowering plants, their chemical composition, use. Vol. 8. (Family Butomaceae – Typhaceae)]. 1994. Saint-Petersburg. 272 p. (In Russian)
19. [Plant Resources of Russia: Wild-growing flowering plants, their component composition and biological activity. Vol. 6. (Family Butomaceae – Typhaceae)]. 2014. Saint-Petersburg. 391 p. (In Russian)
20. Tikhomirova L.I., Bazarnova N.G., Mikushina I.V., Dolganova Z.V. 2015. Farmacol-biochemical study of practical use of some members of the genus *Iris* L. (Overview). — Khimija rastitel'nogo syr'ya. 3: 25–34. (In Russian) <https://doi.org/10.14258/jcprm.201503837>
21. WFO: *Iris lactea* Pall. 2023. <http://www.worldfloraonline.org/taxon/wfo-0000783525>
22. WFO: *Iris oxypetala* Bunge. 2023. <http://www.worldfloraonline.org/taxon/wfo-0000790813>
23. Varlakov M.N. 1933. [East Sayan Expedition NIHFI (Scientific Research Chemical and Pharmaceutical Institute)]. — Khimiko-Farmatsevticheskaya Promyshlennost. 3: 154–157. (In Russian)
24. Varlakov M.N. 1963. [List of plants of Eastern Transbaikalia used in Tibetan medicine]. — In: [Selected Works]. Moscow. P. 122–169. (In Russian)
25. Pryakhina N.I. 1984. [Phytochemical study of milky iris. — *Iris lactea* Pall.: Diss. ... Cand. (Pharmacology) Sci.] Leningrad. 193 p. (In Russian)
26. Minina S.A., Abu Skhela G.R.I., Astakhova T.V., Pryakhina N.I., Zenkevich I.G., Kosman V.M. 1999. Technology of dry extract production from the above-ground part of milk-white *Iris* herbs (*Iris lactea* Pall.) — Pharm. Chem. J. 3(4): 211–213. <https://doi.org/10.1007/BF02509942>
27. Haidav Ts., Altanchimeg B., Varlamova T.S. 1985. [Medicinal plants in Mongolian medicine]. 2nd ed. Ulaanbaatar. 391 p. (In Russian)
28. Li D.-X., Hao X.-G., Zhang S.-K., Wang S.-X., Liu R.-Y., Ma K.-S., Yu S.-P., Jiang H., Guan J.-F. 1981. Antitumor action and toxicity of 6-methoxy-2-delta10'-cis-heptadecenyl-1,4-benzoquinone (irisquinone). — Acta Pharmacol. Sin. 2(2): 131–134. <http://www.chinaphar.com/article/view/4132> (In Chinese with English summary)
29. Han J. 1988. Traditional Chinese medicine and the search for new antineoplastic drugs. — J. Ethnopharmacol. 24(1): 1–17. [https://doi.org/10.1016/0378-8741\(88\)90135-3](https://doi.org/10.1016/0378-8741(88)90135-3)
30. Han R. 1994. Highlight on the studies of anticancer drugs derived from plants in China. — Stem Cells. 12(1): 53–63. <https://doi.org/10.1002/stem.5530120110>
31. Wang X.W. 1999. Irisquinone: Antineoplastic, radiosensitizer. — Drugs Future. 24: 613–617. <https://doi.org/10.1358/dof.1999.024.06.538308>
32. Chen D., Meng Y., Zhu Y., Wu G., Yuan J., Qin M., Xie G. 2018. Qualitative and Quantitative Analysis of C-glycosyl-flavones of *Iris lactea* Leaves by Liquid Chromatography/Tandem Mass Spectrometry. — Molecules. 23(12): 3359. <https://doi.org/10.3390/molecules23123359>
33. Zang J., Xia G., Liu X. 1983. Physicochemical properties of Ma Lin Zi (seed of *Iris pallasii*) oil and identification of its fatty acids. — Chinese traditional and herbal drugs (Zhong cao yao). 14(3): 103–105. (In Chinese)
34. Minina S.A., Pryakhina N.I., Chemesova I.I., Chizhikov D.V. 2008. Medicinal preparation from *Iris lactea* for children. — Khimiko-Farmatsevticheskii Zhurnal. 42(1): 39–41. <http://chem.folium.ru/index.php/chem/article/view/429> (In Russian)
35. Kim J.L., Li H.M., Kim Y.H., Lee Y.J., Shim J.H., Lim S.S., Kang Y.H. 2012. Osteogenic activity of yellow flag iris (*Iris pseudacorus*) extract modulating differentiation of osteoblasts and osteoclasts. — Am. J. Chin. Med. 40(6): 12891305. <https://doi.org/10.1142/S0192415X12500954>
36. Levchuk A.P. 1927. [Hemostatic and uterine agents]. — Bulletin Khimiko-Farmatsevticheskogo NII. 15: 3–79. (In Russian)
37. Chopik V.I., Dudchenko L.G., Krasnova A.N. 1983. [Wild useful plants of Ukraine]. Kyiv. 400 p. (In Russian)
38. Chikov P.S. 1989. [Medicinal plants]. 2nd ed. Moscow. 431 p. (In Russian)
39. Tikhomirova E.A., Sorokina A.A., Bubenchikova V.N., Kostikova E.N., Zhilkina V.Yu., Bessonov V.V. 2020. Chemical Composition and Content of Polysaccharides from the Yellow Iris (*Iris pseudacorus* L.) Rhizomes. — Pharmacogn. J. 12(5): 1012–1018. <https://doi.org/10.5530/pj.2020.12.143>
40. Kashin N.I. 1860. [Home-made medication remedies used by Cis-Argun population, and Cis-Argun folk doctors]. — Bulletin of the Russian Geographical Society. 30(11): 121–145. <https://elib.rgo.ru/safe-view/123456789/218937/1/0L8tMTA0NTZfdmVzdG5pa1JHT18zMGNoXzE4NjAucGRm> (In Russian)
41. Deriker V. 1866. [Collection of folk remedies used by healers in Russia]. Saint Petersburg. 258 p. https://www.travolekar.ru/arch/Deriker_1866.pdf (In Russian)

42. Gammerman A.F., Semichov B.V. 1963. [Dictionary of Tibetan-Latin-Russian names of medicinal plant materials used in Tibetan medicine]. Ulan-Ude. 108 p. (In Russian)
43. Vostrikova G.G., Vostrikov P.A. 1971. [Medicine of the Dersu people]. Khabarovsk. 37 p. (In Russian)
44. Chopra R.N., Nayar S.L., Chopra J.C. 1956. Glossary of Indian medicinal plants. New Delhi. 174 p.
45. Sugawara S. 1937. [Plants of Saghalien]. Toyohara. 490 p. (In Japanese)
46. Efremova N.A. 1967. Medicinal plants of Kamchatka and the Commander Islands. Petropavlovsk-Kamchatsky. 123 p. (In Russian)
47. Semichov B.V. 1958. [On some herbal products of Indo-Tibetan medicine growing in the Buryat-Mongolian ASSR]. — In: [Local studies of the Buryat-Mongolian branch of the Geographical Society of the USSR]. Issue. 2. Ulan-Ude. P. 158–162. (In Russian)
48. Bondarenko A.S., Aizenman B.E., Shvaiger M.O., Mandrik T.P., Bredikhina A.N. 1964. [Antimicrobial activity of some plants]. — [Phytoncides in the national economy]. Kyiv. P. 170–179. (In Russian)
49. Poroshina G.I. 1964. [Influence of some plant antibiotics on lactic acid bacteria]. — [Phytoncides in the national economy]. Kyiv. P. 146–153. (In Russian)
50. Mykhailenko O., Korinek M., Ivanauskas L., Bezruk I., Myhal A., Petrikaitė V., El-Shazly M., Lin G.-H., Lin C.-Y., Yen C.-H., Chen B.-H., Georgiyants V., Hwang T.-L. 2020. Qualitative and Quantitative Analysis of Ukrainian *Iris* Species: A Fresh Look on Their Antioxidant Content and Biological Activities. — *Molecules*. 25(19): 4588.
<https://doi.org/10.3390/molecules25194588>
51. Annenkov N.I. 1878. [Botanical Dictionary]. 2nd ed. Saint Petersburg. 645 p. (In Russian)
52. Petrusheva N.I. 1957. [Fungicidal properties of higher plants]. — *Bulletin Nauchno-tekhnicheskoy Informatsii Nitskotskogo Botanicheskogo Sada*. 2: 41–46. (In Russian)
53. [“Dzeitshar Migzhan” — the monument of Tibetan medicine]. 1985. Novosibirsk. 88 p. (In Russian)
54. [Plants of Tibetan medicine: the pharmacognostic research experience]. 1989. Novosibirsk. 159 p. (In Russian)
55. Hegnauer R. 1963. Chemotaxonomie der Pflanzen. Monocotyledonene. B. 2. 540 p.
<https://doi.org/10.1007/978-3-0348-9387-9>
56. Ishikura N. 1980. Anthocyanins in iris flowers. — *Experientia*. 36(6): 656–657.
<https://doi.org/10.1007/BF01970120>
57. Harborne J.B., Mabry T.J. 1982. The Flavonoids: advances in research. 744 p.
<https://doi.org/10.1007/978-1-4899-2915-0>
58. Yabuya T., Yokota K., Adachi T., Nagatomo T. 1983. High-performance liquid chromatographic separation of anthocyanins in the flowers of *Iris ensata* Thunb. — *Bulletin Faculty of Agriculture University of Miyazaki*. 30(1): 7–13. (In Japanese with English summary)
59. Yabuya T., Nakamura M., Iwashina T., Yamaguchi M., Takehara T. 1997. Anthocyanin-flavone copigmentation in bluish purple flowers of Japanese garden iris (*Iris ensata* Thunb.). — *Euphytica*. 98(3): 163–167.
<https://doi.org/10.1023/A:1003152813333>
60. Kojima K., Gombosurengyin P., Ondogyni P., Begzsurengyin D., Zevgeegyin O., Hatano K., Ogiyara Y. 1997. Flavanones from *Iris tenuifolia*. — *Phytochemistry*. 44(4): 711–714.
[https://doi.org/10.1016/S0031-9422\(96\)00596-1](https://doi.org/10.1016/S0031-9422(96)00596-1)
61. Choudhary M.I., Hareem S., Siddiqui H., Anjum S., Ali S., Atta-Ur-Rahman, Zaidi M.I. 2008. A benzil and isoflavone from *Iris tenuifolia*. — *Phytochemistry*. 69(9): 1880–1885.
<https://doi.org/10.1016/j.phytochem.2008.03.011>
62. Wang H., Cui Y., Zhao C. 2010. Flavonoids of the Genus *Iris* (Iridaceae). — *Mini Rev. Med. Chem.* 10(7): 643–661.
<https://doi.org/10.2174/138955710791384027>
63. Kovalev V.N., Zatylnikova O.A., Kovalev S.V. 2013a. A New Isoflavone from *Iris pseudacorus*. — *Chem. Nat. Compd.* 49(1): 34–35.
<https://doi.org/10.1007/s10600-013-0499-y>
64. Kukula-Koch W., Sienawska E., Widelski J., Urjin O., Głowniak P., Skalicka-Woźniak K. 2015. Major secondary metabolites of *Iris* spp. — *Phytochem. Rev.* 14(1): 51–80.
<https://doi.org/10.1007/s11101-013-9333-1>
65. Wani S.H., Lon S.A., Mustafa M.F., Mir J.I., Qadri R.I., Hassan Q.P. 2017. Evaluation of prominent isoflavonoids of *Iris* plant as futuristic cancer drug components. — *Int. J. Botany Stud.* 2(1): 129–134.
<http://www.botanyjournals.com/archives/2017/vol2/issue1/3-3-45>
66. Zagorulko E.Yu., Ozhigova M.G., Chemesova I.I., Luzhanin V.G. 2018. Quantification of total flavonoid content in the aerial part and in the tincture of *Iris lactea* (Iridaceae). — *Khimija rastitel'nogo syr'ya*. 2: 105–113.
<https://doi.org/> (In Russian)
<https://doi.org/10.14258/jcprm.2018023368>
67. Iwashina T., Mizuno T. 2020. Flavonoids and Xanthones From the Genus *Iris*: Phytochemistry, Relationships with Flower Colors and Taxonomy, and Activities and Function. — *Nat. Prod. Commun.* 15(10): 1934578X2093715.
<https://doi.org/10.1177/1934578X20937151>

68. Bate-Smith E.C. 1968. The phenolic constituents of plants and their taxonomic significance: II. Monocotyledons. — Bot. J. Linn. Soc. 60(383): 325–356.
<https://doi.org/10.1111/j.1095-8339.1968.tb00094.x>
69. Denisova O.A., Glyzin V.I., Patudin A.V., Gavrilenko B.D. 1980. Determination of xanthone glycoside mangiferin content in *Iris*, *Gentiana* and *Hedysarum* species. — Khimiko-Farmatsevticheskii Zhurnal. 14(12): 76–77. (In Russian)
70. Williams C.A., Harborne J.B. 1985. Biflavonoids, Quinones and Xanthenes as Rare Chemical Markers in the Family Iridaceae. — Zeitschrift Für Naturforschung C. 40(5–6): 325–330.
<https://doi.org/10.1515/znc-1985-5-608>
71. Williams C.A., Harborne J.B., Goldblatt P. 1986. Correlations between phenolic patterns and tribal classification in the family Iridaceae. — Phytochemistry. 25(9): 2135–2154.
[https://doi.org/10.1016/0031-9422\(86\)80079-6](https://doi.org/10.1016/0031-9422(86)80079-6)
72. Brooks C.J.W., Watson D.G. 1985. Phytoalexins. — Natural Product Reports. 2(5): 427–459.
<https://doi.org/10.1039/np9850200427>
73. Hoang L., Beneš F., Fenclová M., Kronusová O., Švarcová V., Řehořová K., Baldassarre Švecová E., Vosátka M., Hajšlová J., Kaštanek P., Viktorová J., Ruml T. 2020. Phytochemical Composition and *In Vitro* Biological Activity of *Iris* spp. (Iridaceae): A New Source of Bioactive Constituents for the Inhibition of Oral Bacterial Biofilms. — Antibiotics. 9(7): 403.
<https://doi.org/10.3390/antibiotics9070403>
74. Machalska A., Skalicka-Woźniak K., Widelski J., Głowniak K., Purevsuren G., Oyun Z., Khishgee D., Urjin B. 2008. Screening for phenolic acids in five species of *Iris* collected in Mongolia. — Acta Chromatogr. 20(2): 259–267.
<https://doi.org/10.1556/ACHrom.20.2008.2.10>
75. Pryakhina N.I., Blinova K.F. 1987. Swertiajaponin from *Iris lactea*. — Chem. Nat. Compd. 23(2): 257–258.
<https://doi.org/10.1007/BF00598780>
76. Whaley A.K., Ebrahim W., El-Neketi M., Ancheeva E. 2017. New acetylated flavone C-glycosides from *Iris lactea*. — Tetrahedron Lett. 58(22): 2171–2173.
<https://doi.org/10.1016/j.tetlet.2017.04.080>
77. Chang N., Luo Z., Li D., Song H. 2017. Indigenous Uses and Pharmacological Activity of Traditional Medicinal Plants in Mount Taibai, China. — Evid.-Based Complementary Altern. Med. 1–11.
<https://doi.org/10.1155/2017/8329817>
78. Dragendorff G. 1898. Die Heilpflanzen der verschiedenen Völker und Zeiten. Ihre Anwendung, wesentlichen Bestandteile und Geschichte. Stuttgart. 885 p.
79. Chaurasia O.P., Singh B. 2000. An Ethnobotanical Profile on Cold Desert Flora of Nubra Valley (Ladakh). — Bull. Bot. Surv. India. 42: 81–90.
<https://doi.org/https://nelumbo-bsi.org/index.php/nlmb/article/view/74263/0>
<https://doi.org/10.20324/nelumbo/v42/2000/74263>
80. Haq S.M., Yaqoob U., Calixto E.S., Rahman I.U., Hashem A., Abd Allah E.F., Alakeel M.A., Alqarawi A.A., Abdalla M., Hassan M., Bussmann R.W., Abbasi A.M., Ur Rahman S., Ijaz F. 2021. Plant Resources Utilization among Different Ethnic Groups of Ladakh in Trans-Himalayan Region. — Biology (Basel). 10(9): 827.
<https://doi.org/10.3390/biology10090827>
81. Melnikova T.I. 1994. [Pharmacological study of the crude extract of milky iris (*Iris lactea* Pall.): Abstr. ... Diss. Cand. (Biology) Sci.]. Saint Petersburg. 20 p. (In Russian)
82. Pastushenkov L.V., Barinov V.A., Frolova N.Yu. et al. 1995. [A new immunomodulatory drug – an extract of milky iris]. — In: [Abstracts of the scientific-practical conference dedicated to the 280th anniversary of the First Naval Hospital “Actual Issues of Naval and Clinical Medicine”]. Saint Petersburg. 205 p. (In Russian)
83. Melnikova T.I., Frolova N.Yu. 1996. [Possibilities of using the drug lactir in chronic renal failure]. — In: [Actual problems of creating new drugs: Proc. of the All-Russian scientific conference]. Saint Petersburg. P. 147. (In Russian)
84. Frolova N.Yu., Melnikova T.I., Placen R.I. 1992. [Pharmacological evaluation of the polyextract of *Iris lactea*]. — In: [Abstracts of the 3rd Ukrainian Conference on Medical Botany: Part 1. Kyiv. 139 p. (In Russian)]
85. Gamal R. 1996. [Technology and analysis of tableted dry extract of milky iris (*Iris lactea* Pall.): Abstr. ... Diss. Cand. (Pharmacology) Sci.]. Saint Petersburg. 21 p. (In Russian)
86. Aroyan M.V., Kaukhova I.E., Goncharova S.B. 2017. Determination of merchandising parameters of *Iris lactea* crude sample. — Research results in biomedicine. 3(4): 17–20. (In Russian)
<https://doi.org/10.18413/2313-8955-2017-3-4-17-20>
87. Luzhanin V.G., Whaley A.K., Ponkratova A.O., Zhokhova E.V., Zingalyuk M.A., Pryakhina N.I. 2021. Iris milky-white (*Iris lactea* Pall.) – a promising source of biologically active substances. — Khimija rastitel'nogo syr'ja. 3: 5–17. (In Russian)
<https://doi.org/10.14258/jcprm.2021038890>
88. Astakhova T.V., Pryakhina N.I., Minina S.A., Ismailov G.R. 1994. [On the issue of obtaining and analyzing a dry extract of *Iris lactea* Pall.]. — In: [Solving urgent problems of pharmacy at the present stage: Abstracts of the scientific conference dedicated to the 50th anniversary of the Research Institute of Pharmacy]. Moscow. 240 p. (In Russian)

89. Bychkova N.V., Kalashnikova A.A., Waley A.K., Luzhanin V.G., Kalinina N.M., Shustov E.B., Okovity S.V. 2019. [Study of the immunotropic effects of flavone glycoside embinin by flow cytometry]. — Preventive and Clinical Medicine. 4(73): 77–82. https://profclinmed.szgmu.ru/PIK_4-2019.pdf (In Russian)
90. Astakhova T.V., Pryakhina N.I., Lanina N.E., Zazhigalkina M.V., Chemesova I.I. 2000. [Development of the method for standardizing the syrup with the milk-white iris extract]. — In: [Actual problems of pharmaceutical science and education: Mat. of scientific conf.]. P. 170–171. (In Russian)
91. Lanina N.E. 2003. [Development of technologies for tincture from *Iris lactea* Pall. grass and its standardization: Abstr. ... Diss. Cand. (Pharmacology). Sci.]. Saint Petersburg. 2003. 23 p. (In Russian)
92. Luan Z., Li P., Li D., Meng X., Sun J. 2020. Optimization of Supercritical-CO₂ extraction of *Iris lactea* Seed Oil: Component Analysis and Antioxidant Activity of the Oil. — Ind. Crops Prod. 112553. <https://doi.org/10.1016/j.indcrop.2020.112553>
93. Seki K., Haga K., Kaneko R. 1995. Phenols and a dioxotetrahydrodibenzofuran from seeds of *Iris pallasii*. — Phytochemistry. 38(4): 965–973. [https://doi.org/10.1016/0031-9422\(94\)00731-8](https://doi.org/10.1016/0031-9422(94)00731-8)
94. Wu Sh.-Ch., Chang L., Yang H.-H., Li T.-Y. 1980. Isolation and structural studies of pallasone B and pallasone C. — Yao Hsueh T'ung Pao. 15(8): 43; Chem. Abstrs. 1981. 95: 86202.
95. Zhou J., Xie G., Yan X. 2011. Encyclopedia of traditional Chinese medicines molecular structures, pharmacological activities, natural sources and applications. Vol. 4. Isolated compounds N–S. 636 p. <https://doi.org/10.1007/978-3-642-16779-9>
96. Wu S.J., Yang Q.Z. 1981. Chemical studies on *Iris pallasii* Fisch. var. *chinensis* Fisch. II. The isolation and structure studies of pallasone B and pallasone C. — Acta Chim. Sinica 39: 767–770.
97. Lv H., Yuan Z., Wang X., Wang Z., Suo Y., Wang H. 2015. Rapid separation of three proanthocyanidin dimers from *Iris lactea* Pall. var. *chinensis* (Fisch.) Koidz by high-speed counter-current chromatography with continuous sample load and double-pump balancing mode. — Phytochem. Anal. 26(6): 444–453. <https://doi.org/10.1002/pca.2579>
98. Lv H., Ouyang J., Wang X., Ma X., Suo Y., Wang H. 2015. Separation and Purification of Four Flavan-3-ols From *Iris lactea* Pall. var. *chinensis* (Fisch.) Koidz by High-Speed Counter-Current Chromatography with Flow-Rate Gradient. — J. Liq. Chromatogr. Relat. Technol. 38(15): 1486–1493. <https://doi.org/10.1080/10826076.2015.1063506>
99. Kim H.W., Kim S.S., Kang K.B., Ryu B., Park E., Huh J., Jeon W.K., Chae H.-S., Oh W.K., Kim J., Sung S.H., Chin Y.-W. 2020. Combined MS/MS-NMR Annotation Guided Discovery of *Iris lactea* var. *chinensis* Seed as a Source of Viral Neuraminidase Inhibitory Polyphenols. — Molecules. 25(15): 3383. <https://doi.org/10.3390/molecules25153383>
100. Tie F., Wang J., Liang Y., Zhu S., Wang Z., Li G., Wang H. 2020a. Proanthocyanidins Ameliorated Deficits of Lipid Metabolism in Type 2 Diabetes Mellitus Via Inhibiting Adipogenesis and Improving Mitochondrial Function. — Int. J. Mol. Sci. 21(6): 2029. <https://doi.org/10.3390/ijms21062029>
101. Lv H., Wang H., He Y., Ding C., Wang X., Suo Y. 2015a. Separation and purification of four oligostilbenes from *Iris lactea* Pall. var. *chinensis* (Fisch.) Koidz by high-speed counter-current chromatography. — J. Chromatogr. B. 988: 127–134. <https://doi.org/10.1016/j.jchromb.2015.02.035>
102. Wang X., Jiang B., Lv H., Liang Y., Ma X. 2019. Vitisin B as a novel fatty acid synthase inhibitor induces human breast cancer cells apoptosis. — Am. J. Transl. Res. 11(8): 5096–5104. <https://e-century.us/files/ajtr/11/8/ajtr0091295.pdf>
103. Tie F.F., Luan G.X., Zhou W.N., Wang Z.H., Shi X.B., Li G., Wang H.L. 2018. Effects of the oligostilbenes from *Iris lactea* Pall. var. *chinensis* (Fisch.) Koidz on the adipocytes differentiation of 3T3-L1 cells. — Pharmazie. 73(2): 98–103. <https://doi.org/10.1691/ph.2018.7844>
104. Tie F., Li G., Hu N., Li J., Wang Z., Wang H. 2020b. Oligostilbenes extracts from *Iris lactea* Pall. var. *chinensis* (Fisch.) Koidz improve lipid metabolism in HFD/STZ-induced diabetic mice and inhibit adipogenesis in 3T3-L1 cells. — Biomed. Pharmacother. 131: 110800. <https://doi.org/10.1016/j.biopha.2020.110800>
105. Lv H., Zhou W., Wang X., Wang Z., Suo Y., Wang H. 2016. Extraction and Separation of Vitisin D, Ampelopsin B and cis-Vitisin A from *Iris lactea* Pall. var. *chinensis* (Fisch.) Koidz by Alkaline Extraction-Acid Precipitation and High-Speed Counter-Current Chromatography. — J. Chromatogr. Sci. 54(5): 744–751. <https://doi.org/10.1093/chromsci/bmv249>
106. Tie F.F., Fu Y.Y., Hu N., Chen Z., Wang H.L. 2022. Isolation of oligostilbenes from *Iris lactea* Pall. var. *chinensis* (Fisch.) Koidz and their anti-inflammatory activities. — RSC Adv. 12(51): 32912–32922. <https://doi.org/10.1039/D2RA05176A>
107. Wang X., Qin M.J., Li L., Ye W.C. 2005. Chemical Constituents of Underground Parts of *Iris lactea*. — J. Chi. Pharm. Univ. 36: 517–519. <https://doi.org/10.1002/chin.200505221>

108. Shen W.J., Qin M.J., Shu P., Zhang C.F. 2008. Two new C-glycosylflavones from the leaves of *Iris lactea* var. *chinensis*. — Chinese Chem. Lett. 19: 821–824.
<https://doi.org/10.1016/j.cclet.2008.04.031>
109. Meng Y., Qin M., Qi B., Xie G. 2017. Four new C-glycosylflavones from the leaves of *Iris lactea* Pall. var. *chinensis* (Fisch.) Koidz. — Phytochem. Lett. 22: 33–38.
<https://doi.org/10.1016/j.phytol.2017.08.011>
110. Niu Y.F., Shao Y., Zhao X.H., Wen H.X., Tao Y.D. 2008. [RP-hPLC determination of flavonoids in several flowers]. — Zhongguo Zhong Yao Za Zhi. 33(18): 2102–4. <https://pubmed.ncbi.nlm.nih.gov/19160795/> (In Chinese)
111. Jacobs M.L., Burlage H.M. 1958. Index of plants of North Carolina with reputed medicinal uses. Austin. 322 p.
112. Kimura K., Kimura T. 1970. Medicinal Plants of Japan in Colour. Osaka. 139 p. (In Japanese)
113. Shigemitsu O. 1938. [Medicinal Plant Raw Material]. Tokyo. (In Japanese)
114. Uphof J.C.T. 1968. Dictionary of Economic Plants. Wurzburg. 591 p.
115. Wang S.X., Liu M.F., Yuan L., Zhang W.L., Li D.H. 1986. Effects of irisquinone on cyclic nucleotides in plasma, cancer and other tissues of mice bearing U₁₄ tumor. — Chin. J. Clin. Oncol. 13: 241–243.
116. Li W.M., Wang S.H., Kuang P., Qu B.X., Song H.L., Zhu L.X. 1987. Radiosensitizing effect of an active principle Iq7611 from the seed of *Iris lactea* var. *chinensis*. — Tumor. 7: 97–99.
117. Tang W., Hemm I., Bertram B. 2003. Recent Development of Antitumor Agents from Chinese Herbal Medicines; Part I. Low Molecular Compounds. — Planta Medica. 69(2): 97–108.
<https://doi.org/10.1055/s-2003-37718>
118. Aini H., Ochi H., Iwata M., Okawa A., Koga D., Okazaki M., Sano A., Asou Y. 2012. Procyanidin B3 prevents articular cartilage degeneration and heterotopic cartilage formation in a mouse surgical osteoarthritis model. — PLoS One 7(5): e37728.
<https://doi.org/10.1371/journal.pone.0037728>
119. The Editorial Committee of Chinese Materia Medica, State Administration of Traditional Chinese Medicine. 1999. — Chinese Materia Medica. 22: 271–275.
120. Cui Y.M., Wang H., Liu Q.R., Han M., Lu Y., Zhao C.Q. 2011. Flavans from *Iris tenuifolia* and their effects on β -amyloid aggregation and neural stem cells proliferation *in vitro*. — Bioorg. Med. Chem. Lett. 21(15): 4400–4403.
<https://doi.org/10.1016/j.bmcl.2011.06.039>
121. Orton O., Nadmid S., Paetz C., Dahse H.M., Voigt K., Bartram S., Boland W., Dagvadorj E. 2021. Chromane Derivatives from Underground Parts of *Iris tenuifolia* and Their *In Vitro* Antimicrobial, Cytotoxicity and Antiproliferative Evaluation. — Molecules. 26(21): 6705.
<https://doi.org/10.3390/molecules26216705>
122. Ligaa U., Davaasuren B., Ninjil N. 2005. Medicinal Plants of Mongolia Used in Western and Eastern Medicine. Ulaanbaatar. 134 p. <https://apps.who.int/iris/rest/bitstreams/920761/retrieve>
123. Jalsrai A., Numakawa T., Ooshima Y., Adachi N., Kunugi H. 2014. Phosphatase-Mediated Intracellular Signaling Contributes to Neuroprotection by Flavonoids of *Iris tenuifolia*. — Am. J. Chinese Med. 42(01): 119–130.
<https://doi.org/10.1142/S0192415X14500086>
124. Jalsrai A., Reinhold A., Becker A. 2017. Ethanol *Iris tenuifolia* extract reduces brain damage in a mouse model of cerebral ischaemia. — Phytother. Res. 32(2): 333–339.
<https://doi.org/10.1002/ptr.5981>
125. Pankova P.A. 1949. [Herbaceous plants containing vitamin C]. — In: [Plant raw materials. Proceedings of the Botanical Institute of the Academy of Sciences of the USSR]. 5(2): 292–478. (In Russian)
126. Gubanov I.A., Krylova I.L., Tikhonova V.L. 1976. [Wild useful plants of the USSR]. Moscow. 360 p. (In Russian)
127. Wang Y.Q., Tan J.J., Tan C.H., Jiang S.H., Zhu D.Y. 2003. Halophilols A and B, two new stilbenes from *Iris halophila*. — Planta Med. 69(8): 779–81. PMID:
<https://doi.org/10.1055/s-2003-4279214531035>
128. Kim J.-B., Kim K.-H., Hwang S.-K., Kim Y.-H., Cho K.-J., Hwang Y.-S., Park R.-D., Kim J.-B. 2001. The Composition of Useful Medium Chain Fatty Acids in Eight Plant Species. — J. Appl. Biol. Chem. 44(1): 20–23.
129. Khare C. 2007 Indian Medicinal Plants. New York.
https://doi.org/10.1007/978-0-387-70638-2_809
130. Bhan M.K., George V., Kapoor R., Rishi A.K. 1976. Chemical investigation of local plants: Part I. — Indian J. Chem., Section B: Organic Chemistry Including Medicinal Chemistry. 14(6): 475–476.
131. Iwashina T., Kamenosono K., Yabuya T. 1996a. Isolation and identification of flavonoid and related compounds as co-pigments from the flowers of *Iris ensata*. — J. Jap. Bot. 71: 281–287.
132. Hayashi K. 1940a. Isolation of Ensatin, an Acylated Anthocyanin from the Flowers of Japanese *Iris*. — Proceedings of the Imperial Academy. 16(9): 478–481.
<https://doi.org/10.2183/pjab1912.16.478>
133. Hayashi K. 1940b. On the Anthocyan in the Flowers of Japanese *Iris*. (Preliminary Note.). — Shokubutsugaku Zasshi. 54(637): 23–28.
<https://doi.org/10.15281/jplantres1887.54.23>

134. Hayashi K. 1941. Studien über Anthocyane, VI. Über Ensatin, ein neues Anthocyanin der Blüten von Japanischen Schwertlilien. — Acta Phytochim. 12: 65–82.
135. Hayashi K., Ootani S., Sato K., Tuyama S., Ishikura N. 1978. Chemotaxonomical studies on Japanese irises in relation to the flavonoids in flowers. — Res. Inst. Evolut. Biol. Sci. Rep. 1: 1–16. (In Japanese with English summary)
136. Ishikura N., Yamamoto E. 1978. Anthocyanins in the flowers of Japanese garden iris belonging to “Higo” line. — Ku-mamoto J. Sci.: Biol. 14: 9–15.
137. Karrer W. 1958. Konstitution und Vorkommen der organischen Pflanzenstoffe (exclusive Alkaloids). Basel-Stuttgart. 1207 s. (In German)
<https://doi.org/10.1007/978-3-0348-6808-2>
138. Yabuya T. 1987. High-performance liquid chromatographic analysis of anthocyanins in induced amphidiploids of *Iris laevigata* Fisch. × *I. ensata* Thunb. — Euphytica. 36(2): 381–387.
<https://doi.org/10.1007/BF00041481>
139. Yabuya T., Nakamura M., Yamasaki A. 1994. P-coumaroyl glycosides of cyanidin and peonidin in the flowers of Japanese garden iris, *Iris ensata* Thunb. — Euphytica. 74: 47–50.
<https://doi.org/10.1007/BF00033766>
140. Yabuya T., Imayama T., Shimomura T., Urushihara R., Yamaguchi M. 2001. New types of major anthocyanins detected in Japanese garden iris and its wild forms. — Euphytica. 118: 253–256.
<https://doi.org/10.1023/A:1017562518106>
141. Imayama T., Yabuya T. 2003. Characterization of Anthocyanins in Flowers of Japanese Garden *Iris*, *Iris ensata* Thunb. — Cytologia. 68(2): 205–210.
<https://doi.org/10.1508/cytologia.68.205>
142. Yabuya T. 1991. High-performance liquid chromatographic analysis of anthocyanins in Japanese garden iris and its wild forms. — Euphytica. 52(3): 215–219.
<https://doi.org/10.1007/BF00029398>
143. Kitahara K., Murai Y., Bang S.W., Kitajima J., Iwashina T., Kaneko Y. 2014. Anthocyanins from the Flowers of Nagai Line of Japanese Garden Iris (*Iris ensata*). — Nat. Prod. Commun. 9(2): 201–204.
<https://doi.org/10.1177/1934578X1400900216>
144. Iwashina T. 2003. Flavonoid function and activity to plants and other organisms. — Biol. Sci. Space. 17(1): 24–44.
<https://doi.org/10.2187/bss.17.24>
145. Boltenkov E.V., Rybin V.G., Zarembo E.V. 2005. Flavones from Callus Tissue of *Iris ensata*. — Chem. Nat. Compd. 41(5): 539–541.
<https://doi.org/10.1007/s10600-005-0200-1>
146. Vereshchagin V.I. 1959. [Useful plants of Western Siberia]. Moscow; Leningrad. 348 p. (In Russian)
147. Pozdneev A.M. 1908. [Textbook of Tibetan medicine]. Vol. 1. Saint Petersburg. 425 p. (In Russian)
148. Eom T., Kim J., Lee S., Jeong J. 2013. A herbalogical study on the plants of Iridaceae in Korea. — Korean J. Herbology. 28: 85–93.
<https://doi.org/10.6116/kjh.2013.28.3.85>
149. Ghobadi Pour M., Mirazi N., Seif A. 2019. Treatment of liver and spleen illnesses by herbs: Recommendations of Avicenna’s heritage “Canon of Medicine”. — Avicenna J. Phytomed. 9: 101–116.
150. Kumar S., Pandey S. 2015. An ethnobotanical study of local plants and their medicinal importance in Tons river area, Dehradun, Uttarakhand. — Indian J. Trop. Biodiv. 23(2): 227–231.
[https://tfri.icfre.gov.in/IJTB-Journal/IJTB%2023\(2\)%202015/abstract/IJTB-23\(2\)%202015-\(227-231\).pdf](https://tfri.icfre.gov.in/IJTB-Journal/IJTB%2023(2)%202015/abstract/IJTB-23(2)%202015-(227-231).pdf)
151. Singh K.N., Lal B. 2008. Ethnomedicines used against four common ailments by the tribal communities of Lahaul-Spiti in western Himalaya. — J. Ethnopharmacol. 115(1): 147–159.
<https://doi.org/10.1016/j.jep.2007.09.017>
152. Singh K.N. 2012. Traditional knowledge on ethnobotanical uses of plant biodiversity: a detailed study from the Indian western Himalaya. — Biodiv. Res. Conserv. 28(1): 63–77.
<https://doi.org/10.2478/v10119-012-0028-z>
153. Zaman S., Farrukh H., Muhammad I. 2014. Traditional knowledge on plant resources of Ashezai and Salarzai Valleys, District Buner, Pakistan. — Afr. J. Plant Sci. 8(1): 42–53.
<https://doi.org/10.5897/AJPS12.067>
154. Shuaib M., Hussain F., Rauf A., Jan F., Romman M., Parvez R., Zeb A., Ali S., Abidullah S., Bahadur S., Shah A.A., Azam N., Dilbar S., Begum K., Khan H., Sajjad S., Muhammad I., Shah N.A. 2021. Traditional knowledge about medicinal plant in the remote areas of Wari Tehsil, Dir Upper, Pakistan. — Braz. J. Biol. 83: e246803.
<https://doi.org/10.1590/1519-6984.246803>
155. Kim E.T., Hwang H.S., Lee S.M., Lee S.J., Lee I.D., Lee S.K., Oh da S., Lim J.H., Yoon H.B., Jeong H.Y., Im S.K., Lee S.S. 2016. Effects of Medicinal Herb Extracts on *In vitro* Ruminant Methanogenesis, Microbe Diversity and Fermentation System. — Asian-Australas. J. Anim. Sci. 29(9): 1280–1286.
<https://doi.org/10.5713/ajas.16.0053>

156. Suresh D.K., Ahemad W., Khalid M.S., Aasim S.M. 2010. Anti-hyperglycemic Activity of *Iris ensata* Thunb. Root Extracts in Normal, Glucose Fed and Streptozotocin Induced Diabetic Rats. — Adv. Pharmacol. Toxicol. 11(3): 93–101. <https://www.proquest.com/docview/837356853>
157. Saxena B.P., Srivastava J.B. 1972. Studies on plant extracts with juvenile hormone activity. Effects of *Iris ensata* Thamb. (Iridaceae) on *Dysdercus koenigii* F. (Pyrrhocoridae). — Experientia. 28(1): 112–113. <https://doi.org/10.1007/BF01928298>
158. Spangler J.T., Sample D.J., Fox L.J., Albano J.P., White S.A. 2019. Assessing nitrogen and phosphorus removal potential of five plant species in floating treatment wetlands receiving simulated nursery runoff. — Environ. Sci. Pollut. Res. 26: 5751–5768. <https://doi.org/10.1007/s11356-018-3964-0>
159. Richardson P.M. 1983. The taxonomic significance of C-glycosylxanones in flowering plants. — Biochem. Syst. Ecol. 11(4): 371–375. [https://doi.org/10.1016/0305-1978\(83\)90039-X](https://doi.org/10.1016/0305-1978(83)90039-X)
160. Iwashina T., Ootani S. 1996b. Flavonoids in the flowers of *Iris laevigata* (Iridaceae). — Sci. Rep Res Inst Evolut Biol. 8: 41–48.
161. Iwashina T., Ootani S. 1998. Flavonoids of the genus *Iris*: Structures, distribution and function (review). — Ann. Tsukuba Bot Gard. 17: 147–183. https://www.kahaku.go.jp/research/publication/tsukuba/download/17/ATBG17_147.pdf
162. Marner F.-J., Littek A., Arold R., Seferiadis K., Jaenicke L. 1990. Isolation and structure determination of new spiro-bicyclic triterpenoids from *Iris pseudacorus*. — Liebig's Ann. Chem. 6: 563–567. <https://doi.org/10.1002/jlac.1990199001105>
163. Jaenicke L., Marner F.-J. 1986. The Irones and Their Precursors. Fortschritte Der Chemie Organischer Naturstoffe. — In: Progress in the Chemistry of Organic Natural Products. 1–25. https://doi.org/10.1007/978-3-7091-8888-0_1
164. Zatylnikova O.A., Kovalev V.N., Kovalev S.V. 2013. Composition of *Iris pseudacorus* (Iridaceae) Essential Oil. — Rastitelnye resursy. 49(2): 233–240. <https://elibrary.ru/item.asp?id=18925981> (In Russian)
165. Mikhailenko O.A. 2014. [Comparative analysis of the component composition of the essential oil of fresh and dried rhizomes of yellow iris (*Iris pseudacorus*)]. — In: [New advances in chemistry and chemical technology of plant raw materials: Materials of the VI All-Russian Conference]. Barnaul. P. 210–212. <https://www.elibrary.ru/item.asp?id=22401392&pff=1> (In Russian)
166. Henshaw G.G., Coult D.A., Boulter D. 1962. Organic acids of the rhizome of *Iris pseudacorus*. — Nature. 194: 579–580. <https://doi.org/10.1038/194579b0>
167. Boulter D., Coult D.A., Henshaw G.G. 1963. Some Effects of Gas Concentrations on Metabolism of the Rhizome of *Iris pseudacorus* L. — Physiol. Plant. 16(3): 541–548. <https://doi.org/10.1111/j.1399-3054.1963.tb08331.x>
168. Tarbeeva D.V., Fedoreev S.A., Veselova M.V., Kalinovskii A.I., Gorovoi P.G., Vishchuk O.S., Zadorozhnyi P.A. 2015. Polyphenolic Metabolites from *Iris pseudacorus* Roots. — Chem. Nat. Compd. 51(3): 451–455. <https://doi.org/10.1007/s10600-015-1313-9>
169. Mykhailenko O., Gudžinskis Z., Kovalyov V., Desenko V., Ivanauskas L., Bezruk I., Georgiyants V. 2020. Effect of ecological factors on the accumulation of phenolic compounds in *Iris* species from Latvia, Lithuania and Ukraine. — Phytochem. Anal. 31(5): 545–563. <https://doi.org/10.1002/pca.2918>
170. Mykhailenko O.O., Kovalyov M.V. 2016. Phenolic compounds of the genus *Iris* plants (Iridaceae). — Čes. slov. Farm. 65(2): 70–77. <https://www.prolekare.cz/casopisy/ceska-slovenska-farmacie/2016-2-12/fenolicke-slouceniny-rostlin-rodu-iris-iridaceae-58506>
171. Michalak A., Krauze-Baranowska M., Migas P., Kawiak A., Kokotkiewicz A., Królicka A. 2021. *Iris pseudacorus* as an easily accessible source of antibacterial and cytotoxic compounds. — J. Pharm. Biomed. 195: 113863. <https://doi.org/10.1016/j.jpba.2020.113863>
172. Ali A.A., El-Emary N.A., Darwish F.M. 1993. Studies on the constituents of two *Iris* species. — Bull. Pharm. Sci., Assiut Univ. 16(2): 159–162. <https://doi.org/10.21608/bfsa.1993.70202>
173. Isaev D.I., Kerimov U.B., Kovalyov S.V., Zatylnikova O.O. 2010. [Isoflavones from rhizomes of *Iris imbricata* Lindl. and *Iris pseudacorus* L.] — Pharmacom. 1. 38–42. http://sphu.org/wp-content/uploads/2017/01/Farmacom_1_2010.pdf (In Russian)
174. Isaev D.I., Kerimov U.B., Kovalyov S.V., Zatylnikova O.O. 2009. Xanones from rhizomes of *Iris imbricata* Lindl. and *Iris pseudacorus* L. — Pharmacom. 4: 24–28. http://sphu.org/wp-content/uploads/2017/08/Farmacom_4_2009.pdf (In Russian)
175. Senatore F., Randazzo A., Rigano D. 2003. Pyrrolidin-2-one derivatives from *Iris pseudacorus* (Iridaceae). — Biochem. Syst. Ecol. 31(6): 657–659. [https://doi.org/10.1016/S0305-1978\(02\)00228-4](https://doi.org/10.1016/S0305-1978(02)00228-4)

176. Cardenas M.L., Kickuth R. 1978. Secondary metabolites from macrophytes and their ecochemical function. Part 3. 3-(3-carboxyphenyl)alanine as constituent and excretion product of *Iris pseudacorus*. — *Angew. Bot.* 52: 203–214.
177. Sutherland W.J. 1990. *Iris pseudacorus* L. — *J. Ecol.* 78(3): 833.
<https://doi.org/10.2307/2260902>
178. Zatylnicova O.A., Kovalev S.V. 2009. Study of amino acids and mineral composition of the underground organs of *Iris pseudacorus* L. — *Pharmacom.* 1: 45–47. http://sphu.org/wp-content/uploads/2017/08/Farmacom_1_2009.pdf (In Ukrainian)
179. Tarbeeva D.V. 2016. [Polyphenolic metabolites of *Iris pseudacorus* L. and its cell culture: Abstr. ... Diss. Cand. (Chemistry) Sci.]. Vladivostok. 24 p. (In Russian)
180. Kovalev V.N., Mikhailenko O.A., Kovalev S.V. 2013b. Coumarins of yellow iris *Iris pseudacorus* L. — *Khimija rastitel'no-go syr'ja*. 3: 201–205. (In Russian)
<https://doi.org/10.14258/jcprm.1303201>
181. Hanawa F., Tahara S., Mizutani J. 1991a. Isoflavonoids produced by *Iris pseudacorus* leaves treated with cupric chloride. — *Phytochemistry*. 30(1): 157–163.
[https://doi.org/10.1016/0031-9422\(91\)84117-B](https://doi.org/10.1016/0031-9422(91)84117-B)
182. Hanawa F., Tahara S., Mizutani J. 1991b. Flavonoids produced by *Iris pseudacorus* leaves treated with cupric chloride. — *Phytochemistry*. 30(7): 2197–2198.
[https://doi.org/10.1016/0031-9422\(91\)83613-P](https://doi.org/10.1016/0031-9422(91)83613-P)
183. Sary H.G., Ayoub N.A., Singab A.B., Ahmed A.H., Al-Azizi M.M. 2004. Chemical constituents and molluscicidal activity of *Iris pseudacorus* L. cultivated in Egypt. — *Bull. Pharm. Sci., Assiut University*. 27(1): 161–9.
<https://doi.org/10.21608/bfsa.2004.65438>
184. Bate-Smith E.C., Harborne J.B. 1963. Mangiferin and other glycophenolics in *Iris* species. — *Nature*. Vol. 198. № 4887. P. 1307–1308.
<https://doi.org/10.1038/1981307a0>
185. Zatylnikova O.O., Kovalyov S.V., Osolodchenko T.P., Akhmedov E.Yu. 2012. Investigation of the lipophilic fraction from leaves of yellow iris. — *Вісник фармації*. 3(71): 57–59.
https://dspace.nuph.edu.ua/bitstream/123456789/2055/1/VF2012_3.pdf (In Ukrainian with English and Russian summary)
186. Tarbeeva D.V., Fedoreev S.A., Veselova M.V., Kalinovskii A.I., Gorovoi P.G. 2014. Polyphenolic metabolites from *Iris pseudacorus*. — *Chem. Nat. Compd.* 50(2): 363–365.
<https://doi.org/10.1007/s10600-014-0955-3>
187. Stanley R.G., Linskens H.F. 1974. Pollen: biology, biochemistry and management. 310 p.
<https://doi.org/10.1007/978-3-642-65905-8>
188. O'Connor W.F., Drumm P.J. 1941. Pigments of *Iris pseudacorus*. — *Nature*. 147(3715): 58–59.
<https://doi.org/10.1038/147058b0>
189. Drumm P.J., O'Connor W.F. 1945. The pigments of the yellow iris (*Iris pseudacorus*). — *Biochem.* 39(2): 211–212.
<https://doi.org/10.1042/bj0390211>
190. Steinmetz E.F. 1957. *Codex vegetabilis*. Amsterdam. 136 p.
191. Villeret S. 1974. Sur l'évolution des acides gras majeurs des fleurs d'*Iris pseudacorus* L. — *C.R. Acad. Sci. (Paris)*. 278: 1213–1216. (In French)
192. Seki K., Kaneko R. 1975. Structure of irisquinone: A new quinone from *Iris pseudacorus* L. (Iridaceae). — *Chem. Ind. H.* 8: 349–350.
193. Kato T., Seki K., Kaneko R. 1986. Insoluble monolayers of irisquinone and its related substances at the air/water interface. — *J. Colloid Interface Sci.* 109(1): 77–89.
[https://doi.org/10.1016/0021-9797\(86\)90283-3](https://doi.org/10.1016/0021-9797(86)90283-3)
194. Marner F.-J., Horper W. 1992a. Phenols and Quinones from Seeds of Different *Iris* Species. — *Helvetica Chimica Acta*. 75(5): 1557–1562.
<https://doi.org/10.1002/hlca.19920750509>
195. Zhou J., Xie G., Yan X. 2011. Encyclopedia of traditional Chinese medicines molecular structures, pharmacological activities, natural sources and applications. Vol. 3. Isolated Compounds H–M. 669 p.
<https://doi.org/10.1007/978-3-642-16747-8>
196. Maciejewska-Potapczyk W., Saniewski M., Pełczyńska M. 1974. Sugars in *Iris pseudoacorus* L. seeds. — *Acta Soc. Bot. Pol.* 43: 195–202.
<https://doi.org/10.5586/asbp.1974.018>
197. Lotti G., Aversa V. 1969. [Seed lipids of water plants. — The Italian review of fatty substances. Vol. XLVI. 668–72]. Fisheries Research Board of Canada. Translation Series № 1500. 1970.
<https://waves-vagues.dfo-mpo.gc.ca/library-bibliotheque/139713.pdf> (Transl. form Italian)
198. Tarbeeva D.V., Fedoreyev S.A., Veselova M.V., Kalinovskiy A.I., Seletskaya L.D., Mazurok T.I., Bulgakov V.P. 2013 Polyphenolic compounds from callus cultures of *Iris pseudacorus*. — *Nat. Prod. Commun.* 8(10): 1419–1420.
<https://doi.org/10.1177/1934578X1300801020>

199. Freerksen E., Bönicke R. 1951. Über antibakterielle Prinzipien in höheren Pflanzen I. — Zeitschrift Für Hygiene Und Infektionskrankheiten. 132(5): 417–449.
<https://doi.org/10.1007/BF02149068>
200. Nickell L.G. 1959. Antimicrobial activity of vascular plants. — Economic Botany. 13(4): 281–318.
<https://doi.org/10.1007/BF02885664>
201. Ramtin M., Pahlaviani M.R.M.K., Massiha A., Issazadeh K., Heidari S. 2013. Comparative evaluation of the antibacterial activities of essential oils of *Iris pseudacorus* and *Urtica dioica* native north of Iran. — J. Pure Appl. Microbiol. 7(2): 1065–1070.
<https://microbiologyjournal.org/comparative-evaluation-of-the-antibacterial-activities-of-essential-oils-of-iris-pseudacorus-and-urtica-dioica-native-north-of-iran/>
202. Zatylnikova O.A., Osolodchenko T.P., Kovalev V.N. 2010. Antimicrobial activity of extracts of *Iris pseudacorus* L. — Annals of Mechnikov Institute. 4: 43–47. http://nbuv.gov.ua/UJRN/ami_2010_4_7 (In Ukrainian)
203. Okba M.M., Abdel Baki P.M., Abu-Elghait M., Shehabeldine A.M., El-Sherei M.M., Khaleel A.E., Salem M.A. 2022. UPLC-ESI-MS/MS profiling of the underground parts of common *Iris* species in relation to their anti-virulence activities against *Staphylococcus aureus*. — J. Ethnopharmacol. 282: 114658.
<https://doi.org/10.1016/j.jep.2021.114658>
204. [Wild medicinal plants]. 1965. Minsk. 361 p. (In Russian)
205. Dimkov P. 1978. [Bulgarian folk medicine]: In 3 volumes. Sofia. Vol. 2. 823 p.; 1979. Vol. 3. 678 p. (In Bulgarian)
206. Rollov A.H. 1908. [Wild plants of the Caucasus, their distribution, properties and application]. Tiflis. 599 p.
207. Grintsevich O.M. 1933. [Medicinal plants of the Western region]. — [Materials for the study of the natural productive forces of the western regions]. Smolensk. Iss. 3. P. 19–106. (In Russian)
208. Volkind I.V., Gurevich I.Ya., Uryupov O.Yu. 1976. [Drug handbook for doctors and pharmacists]. Leningrad. 647 p. (In Russian)
209. Yakovlev G.P., Blinova K.F. 1990. [Botanical-pharmacognostic dictionary]. Moscow. 192 p. (In Russian)
210. Frohne D., Pfänder H.J. 1983. Giftpflanzen. 2nd ed. Stuttgart. 137 p.
211. Roth L., Daunderer M., Kormann K. 1984. Giftpflanzen — Pflanzengifte. München. 1032 p.
212. Crişan I., Cantor M. 2017. New perspectives on medicinal properties and uses of *Iris* sp. — Hop Med. Plants. 24: 24–36.
213. Khokhlova N.A., Derkach N.V., Zatylnikova O.A., Kovalev V.N., Volkovoy V.A. 2012. Pharmacological study of *I. pseudocorus* L. — Ukrainian biopharmaceutical journal. 1–2: 42–45.
http://nbuv.gov.ua/UJRN/ubfj_2012_1-2_12 (in Ukrainian)
214. Grossgeim A.A. 1952. [Plant wealth of the Caucasus]. 2nd ed. Moscow. 632 p. (In Russian)
215. Klobukova-Alisova E.N. 1958. [Wild-growing useful and harmful plants of Bashkiria]. In 2 volumes. Moscow; Leningrad. Vol. 1. 218 p.; Vol. 2. 247 p. (In Russian)
216. Geideman T.C., Ivanova B.I., Lyalikov S.I., Nikolaev L.P., Pozharisskaya L.P., Kholodenko B.G., Shkolnikova T.A., Shutov D.A. 1962. [Useful wild plants of Moldova]. Kishinev. 416 p. (In Russian)
217. Bgazhba M.T. 1964. Plant resources of Abkhazia and their use. Sukhumi. 579 p. (In Russian)
218. Ahmed A.H., El Hamshary E.M. 2005. Larvicidal, miracidicidal and cercaricidal activities of the Egyptian plant *Iris pseudacorus*. — J. Egypt. Soc. Parasitol. 35(1): 41–48.
219. Keizer-Vlek H.E., Verdonshot P.F.M., Verdonshot R.C.M., Dekkers D. 2014. The contribution of plant uptake to nutrient removal by floating treatment wetlands. — Ecol. Eng. 73: 684–690.
<https://doi.org/10.1016/j.ecoleng.2014.09.081>
220. Mizuno T., Okuyama Y., Iwashina T. 2018. Flavonoids from *Iris sanguinea* var. *tobataensis* and chemotaxonomic and molecular phylogenetic comparisons with *Iris sanguinea* var. *sanguinea*. — Bull. Natl. Mus. Nat. Sci. Ser. B. 44(3): 135–145. <https://www.kahaku.go.jp/research/researcher/papers/74783.pdf>
221. Hirose R., Kazuta Y., Koga D., Ide A., Yagishita K. 1981. On the Structure of C-Glycosyl Flavones, Flavoyayamenin and Luteoyayamenin, in Petals of *Iris nertshinskia* Loddiges form *albiflora* Honda. — Agric. Biol. Chem. 45(3): 551–555.
<https://doi.org/10.1080/00021369.1981.10864564>
222. Kawase A. 1968. Studies on the Flavonoid of *Iridaceae*. — Agricultural and Biological Chemistry. 32(8): 1028–1032.
<https://doi.org/10.1080/00021369.1968.10859167>
223. Morita N. 1976. Relation of the classification and the component on plant — Application to the chemotaxonomy in connection with flavonoid. — Kagaku kyoiku. 22: 475–483 (In Japanese).
224. Marner F.-J., Longerich I. 1992b. Isolation and Structure Determination of new Iridals from *Iris sibirica* and *Iris versicolor*. — Liebigs Ann. Chem. (3): 269–272.
<https://doi.org/10.1002/jlac.199219920146>
225. Marner F.-J., Simic K., Scholz B., Küster B. 1995. Isolation and Structure Determination of a New Iridal from *Iris sibirica*. — J. Nat. Prod. 58(2): 299–301.
<https://doi.org/10.1021/np50116a026>
226. Bjoroy O., Rayyan S., Fossen T., Kalberg K., Andersen O.M. 2009. C-glycosylanthocyanidins synthesized from C-glycosylflavones. — Phytochemistry. 70(2): 278–287.
<https://doi.org/10.1016/j.phytochem.2008.12.012>

227. Karpitskiy D.A., Bessonova E.A., Kartsova L.A., Tikhomirova L.I. 2022. Development of approach for flavonoid profiling of biotechnological raw materials *Iris sibirica* L. by HPLC with high-resolution tandem mass spectrometry. — *Phytochem. Anal.* 33(6): 869–878.
<https://doi.org/10.1002/pca.3135>
228. Bate-Smith E.C., Swain T. 1965. Recent developments in the chemotaxonomy of flavonoid compounds. — *Lloydia*. 28: 313–331.
229. Andrews P., Hough L., Jones J.K.N. 1953. 238. Mannose-containing polysaccharides. Part III. The polysaccharides in the seeds of *Iris ochroleuca* and *I. sibirica*. — *J. Chem. Soc. (Resumed)*. 1186–1192.
<https://doi.org/10.1039/jr9530001186>
230. Kurentsova G.E. 1941. [Medicinal plants of the Soviet Far East]. — [Proceedings of the Far Eastern mountain-taiga station Voroshilov-Ussuriyskiy]. 4: 131–226. (In Russian)
231. Skalozubov N.L. 1913. [Botanical Dictionary: Popular names of plants in the Tobolsk Governorate]. — *Annuaire du muse de Tobolsk. Livr. XXI. P. 1–87*. <https://www.prlib.ru/item/331768> (In Russian)
232. Krylov P. 1876. [On the folk medicinal plants used in the Perm Governorate]. — [Proceedings of the Imperial Kazan University Society of Naturalists]. 5(2). 131 p.
<https://vital.lib.tsu.ru/vital/access/manager/Repository/vtls:000492012> (In Russian)
233. Grossgeim A.A. 1940. Fam. *Iridaceae* Lindl. — In: [Flora of the Caucasus]. Baku. Vol. 2. P. 200–227. (In Russian)
234. Kaššák P., Kuli M. 2014. Dyeing potential of the *Iris sibirica* flowers. — *Eur. Sci. J. Special Edition* 2: 372–380.
235. Silva J.D., Nascimento M.G., Castro K.N., Andrade I.M. 2015. Ethnobotanical survey of medicinal plants used by the community of Sobradinho, Lus Correia, Piauí, Brazil. — *J. Med. Plant Res.* 9: 872–883.
<https://doi.org/10.5897/JMPR2015.5881>
236. Kovalev V.M., Mykhailenko O.O., Krechun A.V., Osolodchenko T.P. 2017. Antimicrobial activity of extracts of *Iris hungarica* and *Iris sibirica*. — *Annals of Mechnikov Institute*. 2: 57–64.
<https://doi.org/10.5281/zenodo.803878>
237. Hayashi K., Ootani S., Iwashina T. 1989. Comparative analysis of the flavonoid and related compounds occurring in three varieties of *Iris setosa* in the flora of Japan. — *Res. Inst. Evolut. Biol. Sci. Rep.* 6: 30–60. (In Japanese with English summary)
238. Iwashina T., Ootani S. 1995. Polyphenols in *Iris setosa* var. *canadensis* and their chemotaxonomic comparisons with three Japanese varieties. — *Ann. Tsukuba Bot. Gard.* 14: 35–41.
https://www.kahaku.go.jp/research/publication/tsukuba/download/14/ATBG14_35.pdf
239. Ueno N., Takemura E., Hayashi K. 1969. Additional Data for the Paperchromatographic Survey of Anthocyanins in the Flora of Japan (IV). — *Shokubutsugaku Zasshi*. 82(970): 155–161.
<https://doi.org/10.15281/jplantres1887.82.155>
240. Makarov A.A. 1970. [Medicinal plants of Yakutia]. Yakutsk. 179 p. (In Russian)
241. Arifkhodzhaev A.D., Ismailov Z.F. 1980. [*Iris* polysaccharides]. — *Khimija Prirodnikh Soyedineniy*. 6: 822–823. (In Russian)
242. Zakharov A.M., Boriaev K.I. 1968. [Search for plants containing biologically active compounds in the flora of the Middle Asia and Kazakhstan]. — *Rastitelnye resursy*. 4(2): 243–257. (In Russian)
243. Minaeva V.G., Kiseleva A.A., Volkhonskaya T.A. 1973. [Some results of the survey of plants in the Krasnoyarsk Territory for the content of flavonoid substances]. — [Promising useful plants of Siberian flora]. Novosibirsk. P. 170–178. (In Russian)
244. Makarenko A.A. 1897. [Materials on folk medicine of the Uzhur volost, Achinsk district, Yenisei Governorate with an appendix of a collection of folk medicines from the same volost]. Saint Petersburg. 438 p.
<https://elibrary.ru/handle/123456789/229308> (In Russian)
245. Dzhavanshir E.I., Kerimov Y.B. 2005. Study of fatty acids in lipids from four *Iris* varieties. — *Azerb. Eczacil. J.* 5(2): 26–30; *Chem. Abstrs.* 2007. 146: 247424.
246. Bondarenko A.S., Schweiger M.O., Mandrik T.P., Poddubnaya N.M., Skorobogatko T.I., Mozgovaya L.F., Kolesova E.A. 1967. [Antibacterial activity of plants from the botanical gardens of Ashgabat and Dushanbe]. — [Phytoncides, their biological role and significance for medicine and the national economy]. Kyiv. P. 102–106. (In Russian)
247. Mykhailenko O., Kovalyov V., Kovalyov S., Krechun A. 2017. Isoflavonoids from the rhizomes of *Iris hungarica* and antibacterial activity of the dry rhizomes extract. — *Ars. Pharm.* 58: 39–45.
<https://doi.org/10.30827/ars.v58i1.5919>
248. Mykhailenko O., Gudžinskas Z., Romanova S., Orlova T., Kozyra S., Harna S., Volochai V. 2021. The Comparative Analysis of Carboxylic Acid Composition of Four *Iris* Species from Ukraine. — *Chem. Biodivers.* 18(3): e2000969.
<https://doi.org/10.1002/cbdv.202000969>
249. Rybak V., Kerimova G., Korol V. 2021. Investigation of the anabolic activity of dry extracts of *Iris hungarica* leaves and rhizomes on the model of hydrocortisone-induced protein catabolism. — *Čes. slov. Farm.* 70(2): 59–65.
<https://doi.org/10.5817/CSF2021-2-59>
250. Kostić A.Ž., Gašić U.M., Pešić M.B., Stanojević S.P., Barać M.B., Mačukanović-Jocić M.P., Tešić Ž.L. 2019. Phytochemical analysis and total antioxidant capacity of rhizome, above-ground vegetative parts and flower of three *Iris*

- species. — Chem. Biodivers. 16(3): e1800565.
<https://doi.org/10.1002/cbdv.201800565>
251. Yuan Y., Sun Y., Zhao Y., Liu C., Chen X., Li F., Bao J. 2019. Identification of Floral Scent Profiles in Bearded Irises. — Mol. ecul. 24(9): 1773.
<https://doi.org/10.3390/molecules24091773>
252. Zhou J., Xie G., Yan X. 2011. Encyclopedia of traditional Chinese medicines molecular structures, pharmacological activities, natural sources and applications. Vol. 2. Isolated compounds D-G. 525 p.
<https://doi.org/10.1007/978-3-642-16738-6>
253. Singab A.N.B., Ayoub I.M., El-Shazly M., Korinek M., Wu T.-Y., Cheng Y.-B., Chang F.-R., Wu Y.-C. 2016. Shedding the light on Iridaceae: Ethnobotany, phytochemistry and biological activity. — Ind. Crops Prod. 92: 308–335.
<https://doi.org/10.1016/j.indcrop.2016.07.040>
254. Purevsuren Ts. 2004. Bioactive compounds from *Iris potaninii* Maxim. — A thesis submitted for the degree of Doctor of Philosophy in Biology. Ulaanbaatar: National University of Mongolia. 111 p.
255. Medicinal Plants in Mongolia. 2013. Weltgesundheitsorganisation Regional Office for the Western Pacific. 235 p.
256. Ageev V.E. 1989. [Medicinal plants of Adzharia]. Tbilisi. 61 p. (In Russian)
257. Aseeva T.A., Tarmaeva Z.V., Logina O.K., Bukhasheeva T.K., Trifonova G.P., Andreeva L.B. 1987. [Resources of medicinal plants in Buryatia]. — [Plant resources of Transbaikalia and their use]. Ulan-Ude. pp. 41–61. (In Russian)
258. Telyatiev V.V. 1985. [Useful plants of Central Siberia]. Irkutsk. 400 p. (In Russian)

РОД *IRIS* (IRIDACEAE) В РОССИИ: КОМПОНЕНТНЫЙ СОСТАВ, БИОЛОГИЧЕСКАЯ АКТИВНОСТЬ И ПРИМЕНЕНИЕ В ТРАДИЦИОННОЙ МЕДИЦИНЕ

Н. Б. Алексеева^а, М. М. Дорофеева^б, *, Е. А. Варфоломеева^а, Л. Н. Миронова^с

^аБотанический институт им. В.Л. Комарова РАН, г. Санкт-Петербург, Россия

^бСанкт-Петербургский государственный лесотехнический университет им. С.М. Кирова, г. Санкт-Петербург, Россия

^сБотанический сад-институт ДВО РАН, г. Владивосток, Россия

*e-mail: drofa88@mail.ru

Род *Iris* в России представлен 41 видом, 4 вида являются эндемиками, 11 занесены в Красную книгу Российской Федерации и 30 имеют природоохранный статус в различных регионах. В обзоре представлена информация о компонентном составе, биологической активности и лечебных свойствах 16 видов рода *Iris*, произрастающих в России, и об их использовании в народной медицине. Анализ данных показал, что при изучении компонентного состава, биологической активности и медицинского применения, исследователь должен произвести правильную идентификацию видов рода *Iris* L., зная их географическое распространение, экологию, морфологические характеристики. Собранные данные об использовании видов *Iris* в традиционной медицине. Экстракты и выделенные соединения большинства видов ириса обладают болеутоляющими, антиоксидантными, жаропонижающими, противовоспалительными, антибактериальными, противомикробными свойствами. Некоторые виды обладают также цитотоксическим, противоопухолевым, противодиабетическим, противогриппозным, нейтропротекторным, антигипергликемическим, противоаллергическим, антифедантным действием. Большинство выделенных метаболитов представляют собой флавоноиды, изофлавоноиды, антоцианы, терпеноиды, ксантоны, хиноны, фенольные и жирные кислоты. В официальной и традиционной медицине Азии и Европы используются подземные и надземные части растений *Iris aphylla*, *I. lactea*, *I. pseudacorus*, *I. ruthenica*, *I. sanguinea* и др. Статья является итогом многолетнего изучения ирисов в природе и в коллекциях двух ботанических садов, а также обширного изучения литературы, касающейся компонентного состава и лечебных свойств растений.

Ключевые слова: *Iris*, компонентный состав, биологическая активность, фармакологические свойства, народная медицина